37-X666 / serum and clones.

1245

DATE: APR 12 1955 REF:

A. Misc. tests on serum, diluents.
Water in Sutter tap, this lab distilled, or enzyme institute double distilled appears to be suitable diluent (contra "spent broth"), to limit growth. Broth 2.10-20 cells in toto motility, did not flatten unduly and growth was limited. (Unaware contamination with broth of isolation.)

B. 9, 27 ⊕ resp. were completely inhibited by 1: anti-b, 2 anti-i serum. Overnight in B1 (b) large clone of small cells, no motility except a single wriggling cell (plant dead) A12 A.G. In B2, noted clone at center as above but at periphery, net and cords of long cells and filaments (some in serum?) This was also noted later in i-serum.

C 12 & 13 pupm. Freeze degas 12[20] Spot ① to 2[25] DCG picked 13 2[50] - 3[30] Inc. to 5[35] and plate in MGA 60%. Inc. 5[35] PM - 8[30] AM (15 hours).

35 (34 clones + 1 pure swarms) 9 blanks. (probably picked late) - held to 5/31/55 for photo and counting.

Serum inhibition of trails

1246.
1246

DATE: APR 13 1955 REF: degrees freed 12 (~~trails~~ figs)

A. Serum effect microscopic. Bacteria isolated from soil (.23 ml)
B. FA11 (A) × SW666; FA32 (B) × SW666 and put in 1:100 ⁱ, sera in broth. b seems inactivated both very quickly ⁱ i after 30–90 seconds (usually).

Then tried plating. (by c. 345). Harvest c. 350/ml A; 50

(Next middle broke + numbers in B are doubtful.

A – ~~#~~.2 ml samples (est. count is 50 cells/plate). in MGA60 + (1) 0 Plates ~~FA32~~. somewhat messy but profuse T+S.

(6) b 1:100 3 inh. swarms; not all singles.
(5) b 1:1000 5 swarms (3 inh.); 75 singles; No trails.
(4) ~~b~~ i 1:100 Spread contam. But no trails
(3) i 1:1000 6 large 3 small ~~trails~~ swarms (b, i ?) } No tr
why two kinds here ? t

B .1 ml samples (do.) (1–6).

1 Swarms too messy; don't do % T;
2 (MGA) [5 swarms, 56 1's; 7 short trails]; [45; 77 1's;
3 No T or S.
4 No T or S
5 3 inh. S.; No T
6 2 sw (1 spread contam?) No T.

1:1000 is adequate to inhibit trails!

serum inhibits b trails!

abandon i × b system for this study

DATE: APR 13 1955 REF:

37 - X 666

D. SW967 plate in MGA, MGA60 for (minor tails — SPM. in <24 hours but too crowded.

E ca 50 plates — Re. continuously ditto check SW666 MGA60 only — at 2 days, no minor tails same differences: colonies are ●?

F ca 10 plate — 1237A+ for flares — Incubate 5PM - 10 AM. MGA only note singles, not pure?

G control SW673 "

H { 1 MGA
2 MGA 60 → excessive proportion of swarms. (? age of preparation ?. Do not use.

APR 14 1955 Platings of c. 50 initials early in MGA60 + .01 mlce adshow swarms.

	tails
a	+
b	−
c	+
d	+
i	−
k	+
r	+
12	+
B-0	±

9. c, ord would be quite good for further test.

(c. 5 30 PM)

D Replate P 17 A15: excellent development of minor tail 60% MGA; initial only in MGA but will probably show

Cath.

12 49 DIA for Phot. — Refrigerated 10 AM. 5/15. 60% MGA. Remake sister plates

Needs to be done in my own expts.

APR 15 1955
13

	Today	
Serum effects	✓	
Swarm sibs		*
Flares	✓	
Terminals	start	*
E M.		
SW 967 / HA 10	✓ ✓	
Paired subclones for multiplicity of "T" (Use ± T ± ?)		*
Isolations in water?		
Viability pH 4		
Viscosity pH.		
Trails in 40+ ✓ (viability of subclones multi; stiff swarm term)		

APR 15 1955

Notes: Phase var - TH2 ph2 mongh?

Mention to Bruce 6/i — of. bungndof.

Fla^{+} $H.^{a}$

↓

Fla^{-} $H.^{b}$



Serum inhibitors a → x

1248

DATE: APR 14 1955 REF:

	1 APR 15 1955	3	4	5	6	7	8	9

14: n.g.

4/15: Frese degps, FA76a (S. miami 9) → SW666 11AM.
FA37 → x " "

Preliminaries
SW1250, 1252

not tabulated but results indicate that a does not inhibit b trails, part with a/b trails. b inhibits all trails } at 1:10
eswella
1:16

Rescues # 4, 4

3 cells isolated

1131 ? ~~8 cells picked~~. ~~not thio-apt~~.

~~1) pedigree to n_3~~

1) pedigree to n_3. 1/8 gave motile on transfer, found to be mixed +/- Flat: H_1^i (8+:12 - ~~H_1^i~~ H_1^b. ∴ segregation at $n_{4?}$!

2) n_{13} : all −

3) n_{13} : all −

1132 2 cells. followed directly to about n_3:

1) showed Ichem to n_3 ; n_{13}^-

2) 2 sibs both motile, catenated to ~~n_3~~ n_3, n_8 both. n_{13}^-.

11~3

1134 (C3): → 3/22., each then catenated

4/85,

1131) 1) 1 prolapse to n_3. 1/8 ~~full-mutts~~ gave swarm, mixed $F1a^+H_1^i$ $F1a^-H_1^b$. Segr & hyp.

2) 2 cells → n_{13}^-.

n_0 = (Initial chars

1132) 1) 1 chain ~~$n_1 > n_3 < n_{13}$~~ $n_0 - n_3 < n_{13}$

2) 2 chains $(n_0 > n_1 - n_8 < n_{21})(n_0 > n_1 - n_8 < n_{21})$

1133) 1) Swarm : pure (1/4 inviable; 3/4 → H_1^i) but late coal.

1) $n_0 > n_1 - n_1 > n_{13}$. (2 interm.)

1) $n_0 - n_1 > n_1$.

1) $n_0 - n_1 > n_{13}$.





initial only ↑

1134) E3) $n_4 > n_5 \begin{matrix} \nearrow n_{15} \\ \searrow n_{15} \end{matrix} n_{19} > n_{19}$. (3 interm.)

D4) $n_1 > n_4$. 1 interm — n_4 term.

E5) (sw) (T+S) from 1 cell. At n_5 —— n_{17}. chain (F1) F2 = nm. E11 = swarm in nurse

C6)$_1$ ~~n_0~~ $n_0 - n_{23} > n_{33}$.

C4)$_5$ $n_4 > n_5$ ····· $n_5 > n_{10}$ [S isolates $n_{10} > n_{18}$]. 16,50 interm ed.

$n_{16} > n_{26}$

A3)$_1$ $n_1 - n_3 - n_8$? not tested.

B5)$_{10}$ $n_{13} > n_{15}$ ✗ $n_{15} > n_{20}$

④ $n_{19} > n_{26}$

$n_{15} > n_{18}$ ⑥ n_{28}. n_{18}. ~~n_{19}~~

Species 4/15/57

1134 A5 ⎧ $n_2 > n_{12} - n_{12} > n_{28}$ (2nd 3rd)
7 ⎩ $n_2 > n_{12}$ (+7?) ⟨ $n_{12} < n_{19}$; $n_{12} > n_{19}$; n_{40} ~~[illegible]~~.

(100 ?)

(14 mesolations, 6 days [illegible])

~~A4 { $n_1' > n_{14} - n_{16}$. $n_1' > n_{14} - n_{19}$.~~

A4$_2$ $n_1' > n_5$ ⟨ n_{17}. ; n_{20}.

1135 most cool dried. I clare saved 1134) also had trouble drying

B4 ~~F32~~ [illegible] $n_4 > n_{14} - n_{42}$.

[illegible] n_5^2 [illegible] 17 [illegible]

$n_{17} > n_{27}$

$n_6 > n_{14}$ ⟨ $n_{18} > n_{23}$; [illegible] > [illegible] 31 > 439 ; n_{27}.

$n_6^{3+} > n_{16}$ — 19
22 > 27
17 > 30
32.
31.
32.
19 > 32
19 > 29
33 > 38
16
19 > 32
19 > 24
44.

∴ branching ended (case to 1 thereto)
not sooner than n_{19}
not later than n_{16}.

n14 + 13

n27+n15 n45

1138 add chan

n

B21

/

A21 3 C23 bat

4 C33 < 5

22 2 CW

23 1. >13

24 3 C25 13 A32. > 1

25 2 D25 13 A33. 72.

B22 3, 13 A31.

23 3. > 13

24 3. > 10

25 4 E23 13 B34. < 5

C21 . > 4

22 3. > 13

E21 13 B32. > 5

22 13 B33 2, 8 E31 1,



∴ total chan lyctus

n16 —

A21 19.

A22 22. >27

A23 17 > 15. > 30

A24 32

A25 31

B22 32

23 19 > 32

24 19 > 29

25 33 > 38

C21 16.

22 19. > 32

E21 19 > 24

22 44.

$H_1{}^a$ —X

1252

DATE: APR 20 1955 REF:

FA93 (sc 940 4,5,12 a:-) —X Sc 666 . 11AM-12:15 — 12:40 pf
FA37 " "

A) 93 —X 2 35 fuse dyes. 5PM Harvest ~ .5 x 10^3/10 ml.

B) Same dilute 5 22/ml. = 5 22/1 ml .2 ml samples.
B 5 = 1 ml /6.

C) 37-X (sc 1237) 4 02/.4/ml
1. 0.1 ml M & A 60
2 ± "
3 .001 a
4 .001 b

Plate c 3 35 PM
5/31 The results here are not tabulated but can be done.

APR 20 1955 T Repeated. T.

A	1	—	+	C	1 2 3 —	++ >90%
	3	a	+ (reduced in % + extent)		3 a	++ large tiny no in
* some small late fields	4	b	—		4 b.	—* . — dried out before any strands
	5	b	— (Recheck carefully — seems			

F 12/58

Conclusions: effect of anti-a serum is specific, as it works on $H_1{}^a$—X but not on $H_1{}^b$—X. The effect is, however, not complete and may be difficult to measure. Hold plates to photograph. It is possible that early clones are b and late are a or vice versa.

C. 1 2 3 Isolated strains 5/31.

5/31

from 5 plates of B5 and 2B4 looks for any trails or suspicions.

on 1B4 plate only, 3 susp. trails? or clusters.

B4/a

b

c

These may have had early ~~Bs~~ a phenotype with little enough b to swim in presence of anti b.
Plates to DCG to attempt reisolation of these trails.

6/4. DCG found (on replating isolates)

a: gave fuzz 1's and many clusters (ca 100 cols.) (after 3 hour incubation 37 then overnight at 23°. MG photographed.

b: pure 1's

c: mostly swarms – attempt to isolate any 1's (doubtless contain)

post partition of slants

123

DATE: APR 20 1955 APR 21 1955 REF:

Re 1257 pupa.

305 fuse dips

APR 21 1955

1252 pupa.

fuse dips, 4/5 2¹⁰ - 2³⁵ spot 56 (1)

D C B A



A 1-15

D 1-11.

Examine 5:30 Most active without being handled

A 2, 3, 4, 5, 7 B1 B7 B9 C 1 2 3 4 8 14

D 2, 5. Check for circulation & swarm, trail.

5¹⁰ P 22 - Pickerman slides + plate samples

est 10⁴/ml/10ml = 10³/ml plate .02 ml, .1 ml samples in M

activity is posting

DATE: APR 22 1955 REF:

	1	2	3	4	5	6	7	8	9
A	1 (+) wf			—	0				
	2 (1) (1) wf			3	4				
	3 (1) (1) F		v. active	10	4				
	4 (1)			10	4				
	5 (11) F			4	4				
	(1)			1 0 .	4	Research for (+) by trap:			0
	11.			11 —	4				
	10 wf 1.	vs		0 .	4				0
	F + . 1. 1.			6	4				
	11.	daily		14	4				
	11.			1	4				
	15 11.			0 .	3 pd				
B	1 (11) + .	v.c.		4 —	4				
	wf 1.			18	4				
	25 + + . .			1 0 .	3+				
	T T + .			12	4				
	(1) 1.			(1)	4				
	11.			3+ 2	4				
	(+) + .			1	4				
	1. daily			0	4				
	wf F			2/8	4				
	1 (1) not			1	4				
	1 (1) slow			3 —	4				
	30 wf			c. 30	4				
	C2 (1)			swarm	20% - 3+				
	(1)			—	0				
	(11)			4 —	4				
	1.			—	0				
	wf T T			1	0				
	(11)			0 .	(200) 3				
	T 1. daily			20	4				
	40 + .			1	4				
	C11 + . .			50% swarm	4				
	12 (1)			2	3+				
	13 wf			—	0		4/23.		
	15 (1)			3	3+				
	18 (1) 1. . .		"D1" →	40+	(500)		Proved to be swarm now		(10%)
D	1		2	1	3+				
	2		3	swarm	<1	Inv. wf			
	3		4	3	4				
	3 1.		5	0 .	3 typ.				
	5 1.		6	8	4				
	5 1. (1)		7	swarm? 100	3		✓	10% met	
	1. T		8	est. > 50%	4		✓		(over)
	+ .		9	8					
	+		10	10					

Save serums

D1
D7
D8
C2
C11

① from samples ducttly to Processing (plated specific, over)

{ in slabs for whole clone
in MCA tubes for pooled motile
pool Pla⁻ cols. from MCA over to non

Replate

D1 - see infra

D8 - to crowded to count serums ca. 50%

to recover components stocks for later colony tests.

37 [redacted] —x SW666
undivided clones

April 21, 1955

56 (1) isolates, grown to 2^{13} and summarily examined for motiles. Counts are underestimates. No tech losses

NG ~~12~~ 13 Swarms 4 No motiles 6 less than 9 22 10 or more 11

Maximum extimate: 40+. 5 clones were harvested for replating of the intermediate chains.

No.	Est motiles	harv. + nm.	mot left behind	Plate
B3	18	20	4	120 1's ~~many~~; 2 vs T. 800.
B12	28	22 4	10)	11 + 1 vs T 1 cluster
C9	20	28 6	5	1 tail, def. multiple but compact 28 1's.
C1	30	37		4 v.s. tails 000. 31 1's
D1	40	45 10	10)	34 swarms + 13 nm. No tails.

*Replate Residue
8 sw; 92 cols. (own)

Estimates on clones with many chains are therefore moderately low. Some of the nm may have had a motile but this was looked for. However, these drops were not search with a trap owing to shortness of time.

In addition, 4 drops had apparent swarms, but it was difficult to estimate incidence of non-motile elements. Therefore these were blind-picked and plated immediately. (picked to 10 ml, est. 60-70% recovery; plate .02 and .2 ml samples)(This will help evaluate estimate of clone size as 2^{13}.)

Swarm	% mot.	Found	clone size $\log_2$	Plate
C2	20	10%	11	.02 4 sw; 43 1's .2 33 sw — not counted
C11	50	20%	13	.02 4 sw 29 1's .2 5 sw ~ 250 1's
D6	100?	—	10	[No swarms or plated but not counted.]
D8	50+	10%	4	.02 3 1's .2 3 sw; [redacted]

Repeats

The data may be grouped as follows:

mots.	clones	12+ snl
inv	A1 9 11 B2 11 15 C3 5 6 14 D 10 11 ; D8 D5	Initial active:
0	A10 15 B5 C8 D4 A7	6
1	A6 14 B4 7 10 12 C7 10 D2	9
2	B9 C12	2
3	A2 B8 14 C14 D4	5
4	A5 B1 C4	3
m6	A12	1
8	D6 D9	2
"10"	A3 A4 D10	3
11-12	A6 D6	3
14 18 20	B12 B3 C9	3

C15-D1 conf.in tally but not pltg.

D/swarm: Semi 45/55? Flat was removed from DI before plating, the count on residue of 8 swarms: 92 colonies (in $\frac{.92\ ml}{10\ ml}$) is not fair estimate but the ratio must still have $\approx 10\%$. Late segregators?

April 21, 1955. Zur faire trails.

(1). Serum inhibitions. of anti-a, b // b → x b
a → x b
i → x b.

(a) Sm a does not inhibit b → x b, probably specific.
Is b effect specific? Would need a $F1a_1^- H_1^{\pi}$, e.g.
(Input c S. Heidelberg initials?).

(b) Should also be tried on intermediates as early trails might all tend to be H_1^b and agglutinated.

(2). Late branching? Pedigrees + platings of $\begin{cases} n_{1,0} \\ n_{1,3} \end{cases}$ isolates

(3). E-branching < platings of initial sibs
any large trails in sibs to swarms?

(4). Use all segregates H_1^b (a) trail ends
(b) swarm sibs — esp. if $H_1^b P_1^+$.
(c) look for b-resistant trails.

Today: (A) Replate a/. and of a/i → x b.
(B) start (2).

Tonight Dennis notes — summary for (1).

Does b (minnesota) serum also inhibit H_1^a Fla$^+$ → H_1^b trails. 1255

any Fla$^-$ $H_1^{non\,b}$?

DATE: APR 22 1955 REF: 1250

"b"	1	2	3	4	5	6	7	8	9

New Pupps FA10 → SW666 $10^{-10} - 12^{-15} - 12^{-40}$ High.

Freeze trays 240. Hams 441 ~.800/ml.

A. FA10 → SW666 (.1ml) B a → X. (.2ml) ~.102 cells/plate

9A23: trails

			A b → b	B. a → b.
1. —	(Edwards)	—	++	++. 10 swarms, none
2. b minnesota	1/100	b	—	— no trails
3. "	1/1000	b	—	— no trails
4. a	1/100	a	++	± } much the same.
~~5.~~ a	1/1000	a	++	± } appearance more ≅ MOA dilution.

B pupps. may be late, segregated.

Flop: ∴ b, minnesota also inhibits completely. Reaction may be specific for → H_1^b but this cannot be verified unless a Fla$^-$ H_1 can be isolated. (result of 1250B5 plating?).

Search for b trailers. 1252A2 ⑥ ② 1252B2 ⑧ each show max edge (not serum not diff.)

C 2 ○○ } no trailing swarms.
1 ○○○○ }
3 ○○.

1250B5 4 plates

See 1256.

~~5/21 from 1250B4, B5 search carefully.~~

photos of

50

APR 22 1955
APR 23 1955

Tetrazolium- formazan chains preliminary

P22. Prepare stained cultures. Add 1 ml overnight culture to 7 ml broth + TZ .005% .05% Incubate c. 3-4 hours. Also (A) add TZ (1/200 .5%) to 1 ml culture directly.

Best method of preparation appears to be growth for short interval with TZ. Probably only older nongrowing cells willstain.

Refr. to 1 P 23. Test isol. to agar, small liq. drops. Main traouble with agar is confusion from dirt even under oil. Probably better in fluid with a nonmotile culture.

1PM isol. 1) mot. W-2344 to A1

4 PM 3 more to A2

4PM 6 stained W-2802 to small drops near situs C. These were terminally marked. (Z = f gra
Hunt for rare medial marked= 7,9

2 W-1177,(dividing) 9 is terminal, 10 is medial.

of 10 clones found, 3/10 still located "Z". Fate limited - see protocol.

APR 25 1955 APR 28 1955 Try out agar block methods to immobilize. Mostly a.s.f. owing to dirty agar — hard to find divisve cells. Try methods instead. cf. text 1-2yr. Disadvantage of oil chamber is solubility of released Z in oil

But pul. observations above bear out earlier concept

(1) ⬭ → ⬭⬭ as regular mode of division

(2) most Z ~~be~~ unipolar

(3) Z dies out.

DATE: APR 2 5 1955 REF:

Standard system now: SW-940 (FA 93; H_1^a) —x SW 666.

37°

A. Overnight SW666. 1ml + 1 ml .01% TZ broth. Stained 9:00-10:30. Wash and add FA 93 to pellet. Inc 11 AM - 1 PM. (Cf. B); sediment pellet for harvesting motile :

A= cells prestained; 2hours+ phage/

B. 1:1 + FA93 9:30-11AM. Add = vol. TZ broth. Incubate till stained (1PM). Sedime to harvest pellet. (3i/2 hrs. + phage).

B= cells poststained.

Found: many motile initials in each, but almost no motile A were stained (overstai About 2% of motile B were labelled; c. 50% of parent population.

Summary: 28 isolates from B, 2 from A. 5 clones inviable. initially., only one Z (granule) chain died later. Z chains were followed for 4 to 6 fissions. 1 clone gave a swarm (c. 50% motile) = 31B/ E = preponderance of motiles (>10)

8 clones showed E. This appea

If clones are summarized :



as follows :

(+) n	Found	Random expectation
1	4	4
2	3	2
3+	1	{1
Z	0	{1

or, if cell is



The result agrees with random expectation, but possibility of a negative correlation of Z (after n_3 or n_4) should be tested. Original notion if E were polar was expectation of :



if Z had random chance of marking the E or ~E pole initially. However, Z may either inhibit motility if it marks the E pole, or correlation may be one of selection (in terms of age correlation). Further study is made of Z-marked chains, in 2 senses. (over)

APR 25 1955

1254

DATE: 8 P.M.

does this increase 6/56

REF:

	1	app.	3	total	5	6	7	8	9
1.	5 4 4 3 4 z	0 shrinkum. no trans.			4 4/1. 4 3				
10	8 4 4 5 (23) 4 4 4	0 2 0 0 0			4 4 5 5 5 4 4	↓ 1/2. 6 lines 4/; 6 not.	0	c. 0/8	a 1/4
15	1 8.	(all 4 not) 0 only not. in a.				lines 3/8 4'			
18	2 n.	2 not in b.							
19	Never	cool turned taps 9PM							
20									
21	4				4				
22	5 2	0	b. has 0/32		4		(1+,1-)		
23	5 2				5	0/2	. 6.		
24	8		3/8						
25	16 zuf.				4				
26	4				4				
27	8 z								
28	8 z uf.								
29	1 0.								
30	4	0			6				
31	11 2 3	0 6.			3				
32.	4	0			6 (1 var. already)				

Reexamine at 8PM and isolate residual z from pid. # of var z. to eight not dep.

E.T.

conclude 6/56

E.G. 31 was





Reconstruction:

in general, rightmost is Z chain cell.

the others are successively later sibs of it.

e.g.



1254

DATE: APR 26 1955 10:30 AM. REF: 2

④ Age	A	B	C	D	E	F	7	8	+	9	n
4 mos 3 d.	1 0 4	4 4	25 4	0 —		29 22 → 3:05 PM.			2		4
	2 0 3	1 4	1 4	0 4		2					4
	3 0 —	2+				—					0
	4 2 4	15 4	1 4	0 4		18 too close to pools.			1		4
	5 0 4	1 4	4 4			5					3
	6 1	3 4	12			16			2		2
	7 —					—					0
	8 0	3'	0	0 0		3					6
	9 1 4	1 3'	0 3	0 4		2					4
	10 0	1.3	0			23 23 → tube for plating			1		2
	1	18 4	0	2	1	22 18 →			1		5
	0	1	0	1	0	3 2					6
	0	3 3	0	3 0 3	0	3 3					5
	1	38 2	1 4	2 4		37 28 →			1		4
12 30	15 4 4 4 4	3 4	2 4	1 4		50 43 →			3-4		2
1 15	0					—					0
	2 4	3 4	0 4	1 4		6					5
	18 4	2 4	6 4			26 17 → 6 →					2
	19 1. 100,	3.									1
	21 1	0	1	0		2					4
	0	4	0	0		0					4
	1	0 3'	22	4 0	0	23 [illegible]	21 →		2		6
	2	2	0	2+ 0		4					5
	25 3	1				4					1
	1	0 3	0	3 0	4	1	4				4
	42	1	2	2		47	41 →	2+			0
	1	5	0			6		2			0
	30 0	2	0	3 0	0	2 0					5
	0 4	sw 3	0 •			sw no trails	Pick sw later - prev ?				4
	32										
	× ↑	× ↑	× ↑	× ↑		total clones in whole clone.					
	35										

+ is chain in which pertinent motiles appeared.

n = number of passages for which 2 was followed. (not necessarily isolated!)

Save + Recover (1 ml) Pla-sibo in 31B swarm. Plating of sample from the clone gave 44 swarms: 77 colonies, no trails. .01 ml.

50

TZ, methocel; divided clones

1255

DATE: APR 2 8 1955 REF:

Preliminary expts 4/18 showed that 2% Methocel 4000 immobilized cells; they would stay together after fission and form rosettes. Use to test destabilization of Z in a chain, and to isolate after UV or HN. Mix up Methocel in Penassay. The methocel completely immobilizes the bacteria. ?What concentration allows Fla+ to swim?

93→Sc 666 8:30 – 10 AM, add = vol TZ .01% to 12:05. Spin down and refr.

Abandoned.

B) Apr 29. 2 PM stained W-1177 (c. 2 hrs. mixture overnight + TZ broth). Plate out in methocel broth on c.g., over oil. Also isolate a few definite anpmalies.

Conc. Most cells stain unipolar. Rare (5% bipolar, subpolar). Most chains show terminal granule. Occ. cells lyse in random position.

Some exceptions with interstitial granules— probably from subpolar cells.

Need: observations at first division of subpolar and bipolar cells.
should also spot a fair number of unipolar cpntrols.

2 chans

1255

DATE: APR 28 1955 REF:

(+)	1	2	3	4	5	6	7	8	9
A1 A	2-3PM	5PM							
B									
C									
D									
B1 A									
B									
C									
D									
C1 A									
B									
C									
D									
A2 A									
B									
C									
D									
A 3 A									
B2 A									
B									
C									
D									
B3 A									
B									
C									
D									
C2 A									
B									



Exp. n.g. too much slowing up pulses/min added
Method dog or visibility poor. Let go at key
c. 5P27

DATE: REF:

		(−)	C.2 3M spotted	2	3	4	5	6	7	8	9
E1	A	(+)	—								
	B		—								
	C		—								
	D		—								
E2	A		—								
	B		—								
	C		—								
	D		—								
E3	A		—								
	B		—								
	C		—								
	D		—								