U.W.
MICROBIAL
GENETICS

50

Segregation of F'+ from double F'+

30/IV : 1959 REF: cf. P. 42

1 2 3 4 5 6 7 8 9 10 Isolating of various infected strains.

① Purify double F clone on Blac. (See: P 42. 23/IV ; exp.)

② Test individual colonies on fertility × A_1^- and Gal_2^- by spot test and replica plating method (Xyl : 2979 on MXyl.

③ See segregation of F.

Purpose :

① Is Replica plating on MXyl enough to tell whether F_4^+ or F_8^+ or not?

② Is F" segregate or not. If yes, how about the rate.

Results.

① By replica plating on MXyl seeded W2979 on it.

	# of colonies counted	F_4 (+++)	F_8 (+)
WLF4 × W3086 F8			
(1)	46	33	13
(2)	89	66	23
(31)	67	17	50
(34)	48	47	(1) stmil.
(28)	36	36	0

← spontaneous loss? or after [redacted] between two different F's

② By Spot test

Pick typical segregants for expression of F_4 type and F_8 type and test complementality in the fertility on Gal_2 and Xyl_2. (namely Hi for Gal_2 and low for Xyl_2, or Hi for Xyl_2 and Lo for Gal_2.)

See back page.

: Segregation was not observed; ① gives all F_8 type
② gives all F_4 type.

This results can be explained by mutual exclusion during the growth in peptone medium for overnight.

Conclusion : These results seems parallele to the spot test done on 23/IV See P.

Further experiment : Does [redacted] segregant segregate [redacted] the other F'+ after multiplications? How about F_4 & F_8.

In this kind of experiment, I also must replica plate on MGal seeded 2979. To show the reverse relationship for fertility of Gal and Xyl.

Replica

master

$F_2 \rightarrow F_3$ (1)



mMxyl

BGal

$F_2 \rightarrow F_3$

(31)

Replica

master

F_2 F_4



Mxyl

BGal

Further test for

(33) F2type

see front page; compare with the fertility.

sel: Gal_2

sel: A_1

(1) $F_4 \rightarrow F_5$ Same

F_3 type.

not segregate



Gal2

A1

F_4^-

Isolation of (F_4^R) from 7086 F_4

5/IV ; 1959 REF:

Method :

1). Inoculate 7086 F_4 to penassay 5 m/.

→ 2). Shake it overnight on rotator at 37°C. (This makes more efficiently for isolating F^-)

20 plates → 3). Seed 0.2 m/ / plate (on BGal)

4) Irradiate it by U.V. for 10 sec. to each ~~colony~~ plates.

● 5). Incubate it overnight

Ca. 200 colonies per plates → 6). Replica plate it on MGal seeded w 2985 (A_1^-) on it.

of F^- gal: Ca. : 2~3 colonies/plate 7). Pick F^- colony and inoculate it into broth.

8) Retest the fertility a cross times w 2985 on MGal.

9). Test infectability to F_1^{ts} or F_4 or F_8.

Result :

1. # of colonies tested : Ca. 4000
2. # of F^- colonies obtained : 72 (~~abortive~~ only 3 were F^- the other are F^{ts}) recombination-
3. # of F^- colonies obtained : after testing by cross brushing method

see p. 52. and the back page.

4. Rate of F^- obtained

U.W.
MICROBIAL
GENETICS

Isolation of more-infective F' from F4.

Mutant "Rapid infection"

7/14 1959 REF:

• Experimental schedule:

W6 F4 : 10^3 cells /ml F4 + W6 F- 10^8 cells /ml. : 1 ml

incubate overnight

W6 F4 : 10^3 cells /ml F4 + W6 F- 10^8 cells /ml : 1 ml

Repeat 10 times.

• Method: 1st step.

W6 F4 $\times 10^{-2}$ $\times 10^{-2}$: 0.1 ml /5 ml. : Ca. 10^3 /ml.

W6 F- 1 ml + /5 ml. : Ca 10^8 /ml.

+

5 ml fresh penassay.

3 tubes for each.
1 tube for W6 F4
1 tube for W6 F-

1st : 5:30 p.m.

Incubate overnight. — Test fertility by spot test. Save it in cold room.

Use most fertile tube: Repeat again.

2nd W6 F4 $\times 10^{-2}$ $\times 10^{-2}$ $\times 10^{-2}$: 0.1 ml /5 ml Ca. 10^1 /ml 3 tubes each.
control. W6 F4 ① W6 F- ②

3rd Inoculate No 1 into penassay. 5 ml. and incubate it overnight.



Test for immunity to the infection of F_8 (of F_8 $^+$ strains)

9/4 ; 1959

REF: See p 49.

Method



Principle: F_8

$M\ Gal_4 \longrightarrow \times$ ⑤ $M\ Gal_4^+\ F_8^{R?}$

(W3220 F_8)

×

$M^+\ Gal_2^-\ F^-$

(W2979)

o Difficulty is Gal_4 and Gal_2 are not allelic with each others. Therefor, background gives many a black spot. But it may be possible to differentiate between infectable spot and noninfectable spot.

Unsuccessful result



might be 3220 F_8 × 2979 ⑨ M Gal

9/4: Other method testing F_8^R strain. by replica method:

1). W6 ~~W3220~~ F_8 + 3086 F_8^R ⟶ on 5m B Gal

Replica plate on M Gal Streaked W2979 on it.

Ratio of mix. 1 : 1. in pen. → incubate it overnight → seed it on B Gal →

→ Replica plate it on M Gal. (& 2979).

Inoculum size : 10^8/ml.

2	5	#2	W3086 control.
↑	↑	↑	↑
W6 F_8	W6 F_8	W6 F_8	W6 F_8.

Results : All the clones, # 2, # 5, #42, & are infected by F_8, in otherward, these F^- strains are plain F^- in the sense of immunity. But looks more resistant to the infection of F_8. See back page.

(Compair with control)

5 is most promissing.

control

$F_8 \rightarrow$ 3086

×2979 on H Gal.

$F_8 \rightarrow$ #5

×2979 on H Gal.

$F_8 \rightarrow$ #2



×2979 on H Gal.

$F_8 \rightarrow$ #42

×2979 on H Gal.

next step : Infect F_1 to these & strains and see F_2 coming up or not.

(See p. 61)

Test for transfer of Hfr$_3$ to F$^-$.

W3234 (Cy$^+$ Hfr$_3$ Mal$^-$ MOal$^-_2$ Lac$^-_1$) × W1394 (TLB$_1$ F$^-$ S^R)

Test this possibility.

Procedures.

① Purify W3234 on Blac check S^s before use.

② old culture of W3234 1ml + W1394 ~~0.2ml~~ 0.1ml + 5ml phosp.
and incubate it overnight.

③ Streak it on Blac Sm.

④ Replica plate it on MLac seeded W6 – on it. on MLac



Result: negative result See bellow.

Isolation of F_2 from (3000 (Hfr_2 Th) for Adelberg) W4531

13 NOV. 1959

REF: Cf. p 62.

Method:

1. 3000 5 ml overnight culture + 0.1 ml phage W3086 ●
   ↓
   Incubate it overnight at 37°C.
   ↓
   Seed on B Lac Sm.
   ↓
   Replica plate on M Lac seeded W3828 mit.
   ↓
   Look for Recombination + colonies from it.

2. Test Hi-markers with 3000 as a control and infect F_2 to W6−.
   Isolated F_2: 3086 F_2 1, 2, 3, 4, 5; 3000 control 3086 3086 F^+
   × 3828 on M Lac.

   See ~~back page~~.

3. Infect F_2 to W6−, ~~and check~~ and compare W6 F_2 and W3086 F_2 in the pattern of chromosome transfer to F^-.





Infection of F' after killing of F'+ by U.V.

12/4 ; 1959 REF:

1. Wash W6 F4 once by H_2O, and suspend into 1 ml of Water.
   cultural age : Overnight culture.
2. Irradiate it various times, and add to W6−, and incubate them overnight

	U.V. W6 F4	+ W6−	+ 5 ml phenasey broth.
~~[illegible]~~ :	0.1 ml.	1 ml	

- Time of irradiation :

  0, 10 sec, 15, 20, 25,

- Spread 0.1 ml of $\times 10^2$-diluted suspension (over EMB.) : survival test.

Result :

Time of U.V.-irradiation	0	10	Time of exposure (seconds). 15	20	25	60
Infectivity of F4.	−	±	−	+	−	+
survival count: # of survival cells	152×10^6	0	1	2	0	11 / 19
	129×10^6	0	3	2	0	

(11/19 : areas the thickness of the layer irradiated are ca. 2 mm.)

Same number as much as used in infection.) 0.1 ml.



before incubation — after incubation.

Repeat again.: Use thin layer and mix well during and after irradiations.
apply DNase to the sample. Test lysophil. or temp sensitivity

Test ~~of~~ the infectability of ♀ strains obtained from F4 by U.V. irradiation

11/4 ; 1959.

REF: cf. ♀52.

Infectability of ♀ #39, 69, 78 were tested by mixed culture with W6 F4.

W6F4 1ml + ♀ strain 0.2ml + 1ml phosphate

↓

Incubate it overnight at 37°c.

↓

Purify it on Blac Im.

↓

Replica plate then on Hoal seeded W2988 F⁻A₁⁻ on them

Results:

all of these ♀ strains are infectable by F4. (See back page.)

Not immune to the infection of F4.

These ♀ strains are same as F⁻ in the sence of ~~Resist~~ not immune to the infection of F4.

78 may be more resistant than plain F⁻. (3+6)

control.

W6 Fq → × 3086

×2979
on MGal.

W6Fq → #39.

×2979
on MGal.

W6Fq → #69.

×2977
on MGal.

W6 Fq → # 78

×2979
on MGal.

↑
78 maybe promising.

Mix #78 and (Fq) #5, ou #2, ou #42 comes from Fq, and test the fertility.

see p.54

Try all the combinations.

Segregation of F_1 and F_8 from double F. ($F_{1,8}$ / $F_{4,8}$)
(2nd segregation.)

14/V ; 1959 (segregation from 2nd segregants)

REF: Cf. P48, P53.

Test of 1 2 3 4 5 6 7 8 (from master plate / replica plate) 9 10

1. These strain used are obtained from P. 53 - experiment (Replica plate.) of 2nd segregations. # express isolation numbers of p.53. (see p53 front page bellow). Replica plated on MGal seeded W2979 on it.

Results:

From $F_{8,8}$.

1. F_8 type segregants only gives F_8 type only. (Sometimes gives few F^-). This may be reversion.
2. F^- type segregant which comes from $F_{8,1}$ always gives similar F^- type.

From $F_{1,4}$.

3. F_4 type segregant gives F_4 type with very few exceptions.
4. F_1 type segregate few F_4 and many F_1. But some colonies show higher futility (than F^+). They may still have F_4 in it.

[See back page.]

Segregant from $F_{8,1}$. (Example.)

F_8 type #1

(F_1 type) F^- type #1

on MGal ×2979

on MGal X 2979.

○ check if (F^-) arised after infection of F_8 to F^+. It might be the result (obtained) from exclusion between F^+ and F^1 as reported in the exclusion between virulent phage and temperate phage.

○ check. F_4, or F_8 segregated are still gives F^+ or not.



Segregation of F_1 and F_4 from $F_{4,1}$. F_1 type segregant gives only F_1 type after division.



Probably, F_4 may can grow with F_1 without interference. Both F_4 and F_1 can grow within one cell without competition.

F_1 and F_8 are not miscible, but F_1 and F_4 are missible.

F_1 and F_8 exclude with each others. But F_1 and F_4 is compatible in one cell.

(Some colonies ~~assemble~~ assemble to F_4 and some colonies assemble to F_1.)

Interpretation: Both kinds of Fs are spread into the bacterial populations, and they are distributed with different proportion of particles.

Master plate W6 → F_8^- # 2. continued.



×2979 m Kbal

W6 → F_8^-





W6 → F_8^- #2.



Infection of F_1 to F_8^- W6 $\xrightarrow{F_1}$ × F_8^- 3086.

14/V ; 1959. REF: Cf. p. 59. 2, 5, 42. cf. p. 68.

#2, 6 #5, #742.

Principle

1. Mix W6 1ml + F_8^- (M 5 Hist.) 0.1ml + 5 ml phosphate.
2. Incubate it overnight.
3. Purify it on Blue Sm.
4. Replica plate it on M Gal seeded 2979 on it.
5. Pick Hfr colonies and retest by cross streaking, or spotting method.

Result : Seems promissible! Ratio is quite low, but gives Hfr colonies by infection of F_1 to F_8^-.

1. 3086F- is more infectable by F_1 than 3086F_8^-. But both of strains are infectable by F_1. Susceptible to infection of F_1.
2. Infected F_8^- by F_1 gives Hfr colonies. This may be a recombinant of F.

Further experiment:

1. Pick Hfr colonies and test by (cross streaking method) (spot test) (see bellow)
2. Recombine these defective F'.
3. Retest using more larger scale.
4. check it if type male or standard type or F'type.

3849 Trp-
2985 A_1
4297 G_1
2984 P
1394 TL_1
2979 Gal
4589 H-Gal
on M Gal

put into stab. Test infectivity of this F' and test pattern of fertility on various markers.

F_1 × F_8^-



×2979 on M Gal.

H1 for Histidine : This Hfr type mutant seems differ from F_8, however, it is quite useful because H1 transfus character for histidine, it is linked to antigenic characters.
How does it require that character? H1-marker seems shift to Histidine side a little.



W6 → × 3086
control.

Master plate

Replica plate.



×2979
on Minimal

on Black.

compare ratio of inheritance of F1

W6 — × F2 #5.



×2979
Minimal

Comparison of Hfr_2 and F_2 strains.

17/4, 1959

REF: Cf 857

1 conclusive

	2	3	4	5	6	7	8	9	10
B_1	TL	Ara_2	P	Lac	Gal	Try			
	?								
		H1	H1	H1	H1	H1.			

F_2 shows quite same quality as Hfr_2. in transfer of chromosome.



Hfr$_2$ on Mal.

WGF_2 308(F_2 2323 Hfr$_2$ on M Lac B_1

Try Syntrophy.
A_1
G_1
TL
This is mistake: 2979 was spotted on M lac B_1
3824 Lac
3950 A_2



← Syntrophy
TLB_1
← mistake W2979 was spotted by mistake.
on M lac

W6 —
3086 —
W6
3086 +
Test Ara_2 2383
3086 F_2
W6 F_2

Selected H+ Ara^+_2

on M Ara_2 x 2979

Isolation of F_5 from Hfr$_5$ [redacted] (W4537: Hfr$_5$ [redacted])

16/V ; 1959 REF: (W6-, 3086)

Method : Standard method for isolating F' from Hfr strain was used.

1. Replica plated on MXyl seeded 2979 on it, and looked for Hfr colony formit.
2. Test infected F'$_5$ by cross streaking method. (check H1 for Xyl)
3. Infect F_5 to W6-. see back page :
4. Compair fertility-pattern to original Hfr$_5$ (Hfr$_5$ ~ H1 for Th, Xyl, A,)

W4537
Hfr$_5$ → F_5

on D Lac Sm



3086 F_5

1

2

3086+

3086-

×2979
on MXyl.

on MXyl.
×2979.



Ratio.
F^+ F^- F'
17/26 = 65.4% 9/26 = 34.6% 0/26 = 0%

This infection does not give arised ~~(F5)~~ Htr type. Then, what factor was left behind? If more isolates were tested, it may be obtained W6 F5. (Test at least 100).

3086 F_5 —cure by AO.→ 3086 F_5^- ← F_1

↓

? If a factor which controls high fertility was left ~~out~~ back, infection of F_1 to 3086 F_5^- may ~~give~~ results Htr_5 type cell.

Testing for infectivity of $(F_8)F'$. system

17/V ; 1959

REF:

Principle : M^- Gal_4^- F_8 (W3220 F_8) W4534
M^- + F^- (W6 F-) W4354
+ Gal_4^- F^- W3994.

Select on MGal.

Purpose : Does this system work well ?

Result : ok.

Method :
overnight culture : Penassay 5 ml
at 37°C.
↓
spot test,

F- M^- Gal^+ M^- Gal^+ F_8
W3220 W4534
F8 dura

W3994 (Gal_4 F^-)

W6- (M F-)

W3994 + W6-

Use young culture.

MGal

Developed method to look for colony of F.

1. Dilute W4534 into optimum concentration.
2. Spot it on MGal seeded W3994 + W6-, and incubate it 40 hrs at 37°C.
3. Pick colony from the spot, and inoculate it into penassay. (5 ml)
4. Infect the F to 3994 F- (mix culture to it with 3994 and purify it)
5. Test the infectivity by spot test.

U.W.
MICROBIAL
GENETICS

65

Unsuccessful result.

Recombination between F_8^- defective mutants (polysaccharide defect.)

12/V ; 1959.

REF:

P_8^- : Polysaccharide deficient : P_8^+ : wild type.

~~Escherichia~~ strain used ; # 2, 5, K 2, (W3086 F') -.

Combination.

1 2, 5, K2. ; 2 2, 5 ; 3 2, K2 ; K 5, K2 ;

5 2, - ; 6 5, - ; 7 K2, - ; 8 - only



• Experimental conditions.

Age ; Overnight culture of W3086 F_8^- (W3086 F_8^- 2,5, W3086

Ratio : 0.4 ml F_8^- + 1 ml penassay broth.

↓

Incubated overnight at 37°C

↓

Spotted on 2979 & 3994 on MGal. (Control : W3086 W3086 F

Sel: $M^+ Gal_2^-$ or Gal_2^+

• Interpretation for this negative result.

Cell contacts or conjugation does not occur between those defective F_8^{1+} (just like F^-)

After incubation.

Before incubation





Test W4171 (1895 F+ : Novick's F+). W4171 (M) reverted from W1895. sent by Novick.
(Test H1 marker).

21/V., 1959 REF:

purpose: Is it F+ or F'+? Test H1 marker.

method: 1. purify W4171 on Bac. and pick single colony from it.
2. Incubate it overnight.
3. Spot test. on M Gal. Control W6, W6F-.

Result: W4171 shows same fertility pattern as F+ (W6).

Conclusion: W4171 seems not F'.

Summary of W4171
① It carries "Infective F."
② curable by AO treatment.
③ fertility is low as same as F+ on every markers.



on M Gal.

Test (on the infectivity of) F of W4544

20/V 1959 REF: cf. P.61.

Purpose : Does F of W4544 infective? Is it Removable by AO treatment?

Principle : 3086 $F'_{Rec.}$? ——× W6−

(I) Infectivity.

method : 1ml 3086 $F'_{Rec.}$ + 0.1 ml W6− + 5 ml phosay.

↓ Incubate overnight

↓ purify on BMal.

↓ Pick Mal+ and test their fertility in cross × 2979. on MGal.

This strain carries F_1 or F', and is not standard Hfr. Is O3 type Hfr? What was left behind F_0, then?

(II) Treat W4544 by AO. F in W4544 is sensitive to acridine treatment. (see back page)

Result: AO F^-/F^+ 1?/13 ; Actd 2/11 F^-/F^+

Result.

(I) Infectivity W4544 $\xrightarrow{F_1^?}$ × W6−

↓

Result



Rate of infection

33/33 × 100 = 100 %.

on MGal

× 2979

Pick 1 ~ 10 and retest

Result

1. W4544 infect F to F^-.

Effect of UV to infectivity of F'.

22/V. ; 1959 REF:

Experimental conditions.

overnight culture. | W4534 (H- Gal$_4^-$ F$_8$ Hp^s)
W6- (F- H-)
W3994 (F- Gal$_4^-$)

U.V. 6.5 cm.

Spread it onto petri-dish.

1 ml of W4534 → diameter 8 cm.
(Passage culture)

Survival test.

0 min
5 sec
10
15
20
25
30
35

Take one drop of sample after each UV. irradiation.
Spot it on BGal and dMGal streaked a mixture of W3994 + W6- (1:1).

Conclusion:

F' itself or mechanism for infection of F' are sensitive to UV as well as host cell.

↑ on BGal. ↑ stained. Fast green 5% Ac. Ac. Etoh. Petri method

Time of UV irradiation sec



Replica plated on M Lac B, from Blac Sm

Untreated Control.

AO treated



Looks two states.

X3825 on M Lac B,

F8 × F4.

Ref. P42. P50

Recombination of F itself : selection of Hi for (G, or A_2.) Histidine or Hi for both markers.

22/V ; 1959

REF: (F⁻ A, Xyl Mal, Gal Try Pur, Lac P.)

1. Take ● doubly infected compound male. (# 1, 2, 31)
2. Make colony on D Gal
3. Replica plate it on ~~M Gal~~ ~~Tryptophane~~ added ~~W4529.~~ ~~F⁻ Gal⁻~~ W3861 F⁻ A_2^-
4. ~~Look~~ for Hi colonies to ~~Histidine~~ A_2.
5. Pick the colonies ● Hi for ~~histidine~~ A_2. Infect the F' to the other F⁻ ●
6. Confirm stability and uniformity of the progeny. (non-segregational. Heritable.

See p. 42

Replica — Cross hatching.

Ca 500 colonies per plate.

		plate No.	# of Hr reacting
F_{48}	1.	1	—
●		2	—
		3	1.
		4	—
		5	—
		6	2
		7	—
F_{48}	2	8	●—
		9	2
		10	—
F_{48}	31	11	—
		12	●1
		13	—
		14	—
		15	1
F_{88}	C●	16	—
		17	1
		18	1
		19	1
		20	●—

A_2 : F_4

Xyl : F_8.



#10 obtained from control !!
([redacted] of F_8.)
spontaneous mutation ?

3rd Conformation of the Infection of F' to F+.

F_4 —x F_1^+

20/V ;1959 (W6 F_4) (W3086 F^+). REF: cf. P53.

W4518 W4527

1. purify both strains before using.
   Purified on Blac. and inculated into 5ml phosay.
2. Ratio of mix: 1:1 (1ml + 1ml) overnight culture.
   1ml W527 (control)
3. Incubate then overnight.
4. Streak on B Lac Mal Sm.
5. Replica plate it on M Gal seeded W2985 on it. $R^- A_1^-$
6. Pick Single colony of $F_{4.1}$ and suspend it into phosay. Streak it on Blac Sm.
7. Replica plate it on M Gal seeded W2985 ($A_1^- F^-$).

Results and conclusions.

1. $F_{4.1}^+$ is infectable by F_4. and gives $F_{4.1}$ and F_0 probably $\pm F_r$
2. The percentage of infectivity of F4 to F_1^+ is ca. 1 %.

on Mgal.
x2979

6/669 x 100 = 0.90 %

ca. 1 %

Scheme. $F_4 \times F_1^+$

$\downarrow$

F_{41}^+

$\swarrow \quad \searrow$

$F_4^+ \quad F_1^+$

Segregant F_1 type.



Double F type F_{41}. # 1

W6 $\xrightarrow{F_1}$ × 3086 F_4^d Control $\frac{F_1}{W6}$ × 3086.

W6 $\xrightarrow{F_1}$ × 3086 F_8^d

20/4 ; 1959 REF: Cf. P.61

method : Same as P.61.

Strains : 3086 F_4^d #78 (see p.59), 3086 F_8^d #2.
W6 : F_1 donor cultural age : overnight culture.

Control W6 ⟶ 3086.

Replica plate on MG+Gal Seeded W2979 (Gal⁻ F⁻)
(Select M Gal⁻.)

Replica plate. Retest the Hfr colonies by Cross-lining method

	No of colonies tested	Hfr colonies (Hfr for Gal.)	Hfr clones	
F_1 W6 × F⁻ 3086	286	1	1	1
	255	0		
	334	0		
	269	0		
	256	0		
	232	0		
	Σ 1632	1		
F_1 W6 × F_8^-	88	0		
	74	0		
	91	0		
	103	0		
	94	0		
	450	0		
F_1 W6 × F_4^-	185	0		
	206	0		
	240	2	1	3
	158	0		
	168	2	1	
	162	1	1	
	Σ 1119	5		

Retest by cross-lining method

Conclusion :

1. Test infectivity and accessibility to UV-treatment.
2. Remove F and reinfect F_1 to it. See what happens.

1. Even plain F⁻ gives Hfr colony by infection of F_1. This is unexpected result.

But, this suggest that F⁻ can mutate into F_3 type female by itself, or F_1 can mutate into F' by spontaneous mutations, and become F'⁺ or Hfr.

2. All these Hfrs are not F4 type. This means 3086 F_4^d has no F4 character in there.



Retest on MG+Gal 2, 4, 5, 6, Hfr marker by spot test (see back page)

~~W3898~~ → W3086
W3924

23/4 : 1959. REF:

Hfr x Lys⁺ Thr⁺ Leu⁺ V₁ P100.
Jacob's Hfr.

Method : Just same as standard method.

Replica plated on MAra plates seeded W2979 on them.
[Meth - Ara₂ selection.]



BLac Sm.



x 2979.
MAra



MAra
x 2979