151

Last trial on the killing of cells without loss of mating capacity.

17/xo 1957 REF:

5 ml overnight culture of W6 Fg. in Pen

↓

Centrifuge, discard the supernate, resuspend in 1 ml of H_2O.

↓

distribute the 0.2 ml of W6 Fg into dupont soln. {sodium lauryl sulphate 1:100 PR

↓

Incubate it for 2 hrs. at 37°C.

↓

Wash it once and spot them on MGal x 4573 & on BGal

Conc. of dupont.	survival	fertility.
1 %	+	+
0.5 %	+	+
0.1 %	+	+
control. 0 %	+	+





Conclusion: Use much conc. of dupont. (20 %)

U.W.
MICROBIAL
GENETICS

152

2nd trails on the killing cells, without removal of mating capacity.
(duponol)

17/IX ; 1959. REF:

1 2 3 4 5 6 7 8 9 10

1. W6 F_8 overnight culture 8 ml / tube. 4 tubes.

Centrifuge - Wash once - put 4 tubes together - resuspend them into 1 ml H2O.

3. Add 0.2 ml of the Bacterial suspension into duponol soln. 0 % 10 %
15 % 20 % and incubate them for 4 hrs. at 37° C.
3:00 PM ~ 7 PM.

4. Test survivals & fertility of these treated and untreated W6 F_8.
on B Gal and B M Gal x 4573.

20 % 15 % 10 % 0

1/19 1 ← 2 days old.

Penicillin grown.

Exp. I. 4573 + duponol-lysed F_8^+ (with Duponol) → incubate at 37°c for 30'

0.2 ml original 0.5 ml
0.2 ml 1/10 diluted 1 ml

→ wash once → ~~spot~~ the 0.1 ml on M Gal. (7:30 PM ~ 8:00 PM,
(centrifuge for 5 min.)

Result.

	survival	Recombinant
10 %	2×10^3	$> 2 \times 10^3$
15 %	"	"

Transfer of Lac.

Exp II. dilute the duponol-lysed material (10^{-1}) → spot on MLac on which W4573 was streaked.
non-diluted material (10^0) → "

Untreated control. 0 10 % 15 % 20 %

~~Centrifuge and resuspend in 1 ml H2O and spot on MLac x 4573~~ 1st tube of

Conclusion: ① Killed cells with duponol does not transfer Lac to F^-.

② However, the treated material shows recombinating activity in transfering Gal as same numbers of survival cells.

x 4573

1/10 dil on MLac 1 mail into

Next step: ● Make exact experiment.

4:30

4297 4296 2985 4539 4627 4354 4295
G T A_1 H Trg M L

1
2
3
4
5

blank #295 #354 4627 4539 2985 4296 4297

on MGluc.



4297 4296 2985 4539 4627 4354 4295 blank

on MGluc.

Continued from 153a.

F status of segregants from ♀ 3747 × 4574.

4574 : F^- P Ara$_2$ Mal$_8$ Mtl Ara$_2$ Xyl$_2$ Gal$_2$ SR. primary exconjugants of (cf. P)

24/II ; 196 REF:

Exp. I This experiment does not work well. The fertility was so low in this time. only one Lac$^+$ was tested.

25' : at 37°C. 1 recombinants (Lac$^+$) — purified on BLac Sm — Pick Lac$^+$ & Lac$^-$ onto in incubator. BLac Sm — Replicate on DO + 4506 (F^- pur$^-$)

(Lac$^-$ colonies (4574) are so crowded around the Lac$^+$, therefore, Lac$^-$ F^-(4574) can be contaminated with segregants)

#1 Lac$^+$.	# of Lac$^+$ tested	sex	# of Lac$^-$ tested	sex
	15	F_{13} : 15 F^- : 0	24	F^- : 21 F_{13} : 1 ← Save this F^+ : 2 (or F_{17}) recheck!

#of Lac$^-$ Lac$^+$ 876 : 1

See back page Ia, & Ib.

Exp. II Exp. does work beautifully.

25' : at 37°C : in incubator room : on rotator.

Lac$^+$. tester : 4506 on DO.

① Mix 4hrs exponentially growing and culture of 3747 (10) & 4574 (1) Shake it on rotator for 25' at 37°C. Chill & Then blend the mix, dilute it into 10^{-2}. 0.1 ml. plate on BLac Sm.

② pick well separated Lac$^+$.

③ purify them on BLac Sm. Pick Lac$^+$ segregants & Lac$^-$ segregants. and test sex by Cross-brushing method on DO. × 4506 (See back page II.

	Lac$^+$ tested	sex	Lac$^-$ tested	sex.
#1	8	F_{13} : 8	10	F^+ : 10
#2	10	F_{13} : 10	1	F^+ : 1
#3	4	F_{13} : 4	1	F^+ : 1
#4	8	F_{13} : 8	—	—
#5	6	F_{13} : 6	—	—
#6	7	F_{13} : 7	—	—
#7	8	F_{13} : 8	—	—
#8	6	F_{13} : 6	—	—
#9	5	F_{13} : 5	—	—
#10	6	F_{13} : 6		
Σ	58	58	12	12

% of F_{13} of Lac$^+$ segregats : 100. % of F^+ of Lac$^-$ segregat : 100.

• F^+ means as follows :
1. Not F^- from the cross × F^-
2. Not F' by low fertility.
3. Same fertile as F^+ in cross × F^-.

Fertility of F^+ & Infectivity of F^+ should be checked.

• 3 colonies out of 10 showed segregation of Lac$^-$ #1, 2.

— : No Lac$^-$ colonies can be picked after streak on BLac Sm.

Conclusion : 1. Lac$^+$ becomes F_{13}. suggesting close linkage between Lac & F_{13}.

2. Lac$^-$ becomes F^+. (These F^+ (presumably?) should be checked. in its purity, and low fertility & infectivity for many months) → These are really F^+! see back page.

Further experiment : Pick 3 colonies from #1, #2, #3. and test compatibility of each 3. 1, 11, 12.6 (see back page II)

Speculations : 1.) F^{1+} (cell) carries wild type F, or at least F' which can distribute F into their progeny after mating in separate way from F'. This may be interpreted by 2 alternatives : 1. F'^+ carries wild type F ; 2. F' is easily separate from F and the newly formed F will spread over the progeny.

2) Hfr may be defective F^+. It carries defective F by recombination between F and bacterial chromosome.

Ia:

Lac⁻ ⟷ Lac⁺ on DO ×4506

Ib. · Retest fertility of all Lac⁺ and 2 fertile Lac⁻ and sterile Lac⁻.

Make cross-streaking.

on DO ×4506 Lac⁻ / Lac⁺

II. ~~Tested by~~ Sex of Lac⁻ segregants from primary zygote

Test of fertility of Lac⁻ segregate from Primary zygote



× 4506 on DO. Pm F⁻

on DO.

Test for dual structure of F'-ness.

(Is F' defective? Usual F'^{+}_{REF} contains standard F.)

10/II ; 1960

1 2 ~~3 hrs~~ 3.5 hrs 4 5 6 7 8 9 10

Cultural age : 11:30 ~ 15:00 Inoculum size : 0.2 ml / 10 ml pen.
Shaken on rotator at 37°C. Overnight.
Pen. grown culture

Principle. : Does F' segregate out F after infection of F_{13} from F_{13}^{+} (3747) to F^{-} (4574)? F^{-} Lac$_{85}$ P Ara$_2$ Gal$_2$ Xyl$_2$ Mtl, Mal, S^R.
tester { Lac$_{85}$ Pur^{-}. F^{-} φ^{+} F_{13} V_6^R M
W4642
S^R W4647

Experimental conditions to set it up. :

Use excess F^{0+} (10 times more). Blend the mix with mixer.
(Add 1/ml F' to 10 ml F^{-})
20':10, 25':10, 30':10
Seed on Blac Sm. pick Lac$_V$. Purify each Lac$_V$ on Blac Sm.
Restreak the Lac$_V$ on Blac Sm. again.
Pick Lac^{-} and Lac^{+} from the segregants. Test sex selecting X^{+}
10 10
on M glucose. seeded W6- on it.

Time of Interruption : 0' 10' 15' 20' 25' 30'

Speed of blending 70 : 1 min. Dilute the mix into 5×10^{-4}

Result :

1). Timing experiment : at 34°C, standing in water bath.

Time of blending after mix	0'	10'	15'	20'	25'	30'
# of Lac^{+} S^R	0, 0, 0	0, 0, 0	0, 1, 0	11, 9, 12	38, 35, 23	155, 129, 121
Σ Lac^{+} S^R	0	0	1	32	96	405



Lac^{+} transfer by F_{13} (3747) in Pen. at 34°C.

Conclusion:
↑ Use 25' and try more diluted 1/10 sample.
Use 10 plates for only 25'.

2). Repeat this experiment at 37°C. in incubator room. in pen. 10 ml. on rotator.
- Cultural age : 13:00 ~ 16:00 at 37°C on rotator
- Inoculum size : 0.5 ml / 10 ml of pen. (overnight grown)
- Mix 10 : 3747 with ~~4574~~. 1 ml : 4574.
- Select on Blac Sm. 10^4 : 0.1, 0.5
4:20 ~ 4:45

Result :

Lac^{-} / Lac^{+}
428 : 4
967 : 4
419 : 3
429 : 4
454 : 3

Streak Lac$_V$ on Blac Sm. and pick Lac^{+} & Lac^{-} colonies & test sex-compatibility.

continue to 153b

This data in in 153b.

U.W.
MICROBIAL
GENETICS

154

Experiments to show whether W4630 (Lac_2) homogenote or heterogenote?

22/II. 1960

REF: Lac_2 F^+ : 2242
S^R Lac_2 F^- : 3112.

1 2 3 4 5 6 7 8 9 10

Experimental design:

1. Pick independent Lac_2^+ reversions and purify them on BLac. (Pick 20.)
2. Treat those Lac^+ with AO, and see are there any Lac^- segregate or not.

As a control, use Lac^+ revertant from 3112. (This strain should not segregate Lac^- after AO-treatment.) also F^+ exp 3112 Lac_2^-/Lac^+. (this segregate Lac^- by AO)

3. Check Lac^- segregant for Lac_2-ness.

Points expected for this expts.

1. If it is homozygote for Lac_2, all the Lac^+ does not produce Lac_2^-.
2. If it is homozygote for Lac_2, three possibilities can be expected.
   a. Lac_2^- will be segregated after AO treatment from $Lac_2^-/Ex\ Lac_2^+$ (exogenotic revertants).
   b. Lac_2^- will not be segregated after AO-treatment from Lac_2^+/Lac^- Ex (endogenotic revertants).
   c. Suppressor mutation in outside of F_{13} will not segregate Lac^-.

Experiments:

1. Purify W4630 on EMB Lac Sm. and pick 20 Lac$^-$ single colonies. Suspend these colonies in 1 ml distilled water and spread on BLac separately. Test all F_{13} Lac_2^- by cross-streaking with 4573 (Lac_{95} F^-) on MLac. ~~Inoculate the colony in 8 ml nut. broth.~~ (See back page 1)
2. ~~Inoculate 0.2 ml of the culture on MLac and~~ Incubate them for 24 hrs. at 37°C. Pick Lac^+ from each Lac^- 4630 colonies.
3. purify the Lac^+ colonies on BLac. 20 colonies.
4. Treat these revertants with AO. exponentially growing cell.

Results:

Medium : Nutrient broth : pH. 7.6 : AO : 20γ/ml : 37°C. overnight @100 rpm-Amco

Inoculum size : one loop of 10^7 cells/ml 4630 Lac^+ revertants / 1 ml of medium. 20γ/ml of AO.

Isolation No. of 4630 Lac^+ Revertants.	Lac ν or + before AO	Lac ν or + after AO	Isolation No. of 4630 Lac^+ revertants	Lac ν or + before AO	Lac ν or + after AO
1	No	No	16	No	No
2	No	Segregate	17	No	No
3	No	Segregate	18	No	No
4	No	No	19	No	Segregate
5	No	No	20	No	No
6	No	Segregate	control.		
7	No	No	3112 Lac^+	No	No
8	No	No	3112 $Lac_2/Ex\ Lac^+$	Seg	Seg
9	No	No			
10	No	No			
11	No	No		Seg: 2	Seg: 6
12	No	No	Σ Lac ν / total No. tested	No: 18	No: 14
13	Segregate	Segregate			
14	Segregate	Segregate			
15	No	No			

Streak the treated & untreated cells on EMB Lac. and see are there any Lac^- segregation or not.

6. Restreak from 12, 13, 14, 15, 18, 19 (Single colonies and make sure.) 5. Pick Lac^- segregate and test the Lac marker by cross ×3747 × Lac_2F_{13} on MLac

4630
Isolated from streaking
on Phac Sm



X 4573
m M Lac

segregants from Lac mutants of 4630



V4630

X3747
on M Lac



Compound structure of F' in 3747.
Testing for segregation of F' and F from primary exconjugate.

25/II ; 1960 REF:

Cultural age : exponentially growing culture in 10ml plain broth on rotator. (Inoculum size : 1ml/10ml plain)
4:30 ~ 6:30

Mix them : 3747 : 10ml + 4574 1ml.

↓

Keep them on rotator for 25'

↓

Dilute and chill them in distilled water, then blend the mix.

↓

Dilute it into 10^{-1} : 0.5 : 10^{-2} : $10^{-2} \times 5$: 0.1/plate

↓

Inoculate the 0.1ml on EMB Lac 3m.

Result :

	# of Lac⁻	# of Lac⁺
	367	208
	319	196.

Sex

	Lac⁺		Lac⁻	
	# of Lac⁺	Sex	# of Lac⁻	sex
# 1	3	F13	7	F⁻ 5 ; F⁺ 2.
# 2.	10	♂ 10	5	F⁺ 5
3.	8	" 8	9	F⁻ 9
4	3	" 3	9	F⁻ 9
5	5	" 5	9	F⁻ 9
6	4	" 4	9	F⁻ 9
7	2	" 2	8	F⁻ 8
8	2	" 2	10	F⁻ 10
9	6	" 6	9	F⁻ 9

Testa 4506
Media selected } M6 line.
Spit.

[redacted] of colonies which segregate
: Lac⁻ P⁺ : 2
: Lac⁻ P⁻ : 7

Conclusion : ① F' segregate F to their progenies after mating. But it seems not always occurs. Some F' does but the others are not.
② Fertility of Lac⁺ varies, some shows high the others shows lo.
There seems no distinct difference.
Only a General principle is Lac⁺ is always male.

Test Infectivity of F and F'.

U.W.
MICROBIAL
GENETICS

Transformation of Gal with λ helper.
(Dr. Keien's system)

22/III. ; 1960 REF:

Helper : 434 hg Cs 21 10^{12}/ml

dilute 1, 2, in 2MNacl. 10, 10

Recipient :

W3104 (434 hg / T_1) infected with λ. :

grown in. Pmed { .02 M KPO_4 pH. 7.0 ; Conc. of Bact. 4×10^9/ml.
.001 $MgSO_4$
.001 $(NH_4)_2SO_4$
10^{-6}M Fe
0.5 mg/ml glucose.

OD: 1.78

DNA : Extract DNA from W4687 (c.f. 19/III/60). : Dialyzed at 5°C for 3 hr vs. 2l. vs. .15M NaCl .01 Tris pH. 7.4 ID IID.

Reaction Mix :-

0.1 ml DNase (control).
0.1 ml DNA. (200 μg . 2 μg) W4687
0.1 ml Bacteria.

Keep the mix at 30°C for 1 hr.

		on TTC Plaque	on TTC Gal	(Dilution)	on B Lac Sm	DNase 30 μg	DNA	# of Gal+/ from replicas (on BGal)
1				(1000)	0	—	I 0.1 ml	±48
2	2	341	198	(500) ml	0	—	I 1/10 0.1 ml	+2/+6 ; ±4
3	3			(10) ml	0	—	II 0.1 ml	+0, -6 ±2
4	4	73	41	(10) ml	0	—	II 1/10 0.1 ml	+3, -5
5	5	(423×4) 1692		(500) thin	0	—	I D 0.1	+0, -10
6	6	(423×2) 846	211	(500) thin	0	—	I D 1/10 0.1	+0, -17
7	7	58		(2000) thin	0	—	II D 0.1	+0, -17
8	8	44		(1000) thin	0	—	II D 1/10 0.1	+0, -18
9	9	0	660	(500) ml	0	0.1	I D 0.1	+1, -14
10	10	136		1/500 thin	0	—	λ dg DNA 1/6 10^4 0.1	+15, -0
11	11	0	73	100 ml	0	—	—	remeasure +0, -11 control.
12	12	0	0	0	0	—	I 0.1	
13	13	0	0	0	0	—	II 0.1	
14	14	0	0	0	0	—	I D 0.1	
15	15	0	0	0	0	—	II D 0.1	

F8

W3104 on 2/4.

Gal+ ↑ W3104

Gal+ ↑ W3104

λ dg. 0

λ dg

Gal+ ↑ (T3110 Sp)

Conclusion : 1. DNA extracted from 4687 shows plaque forming activity on W3104. (434 hg/T_1)
2. Some colonies which form TTC+ shows Gal+ quality on BGal.
3. Such activity would be loose by treatment of DNase.



on B Gal.



on B Gal.

U.W.
MICROBIAL
GENETICS

Extraction of DNA from Gal-heterogenote W3094D. (4/ex 2)

23/III 1960 REF:

[redacted]-source : (from W3094 derived strain (4pt Gal+ F⁻)) 4/ex 2. (Made by HFT-Gal$_2$ transduction)

21/III

1. Shake it on shaker (2.5 l in 4 l flask.) at 37°C for overnight.

22/III

2. C.f.g. collect, and wash the cells with 0.14 M NaCl 3 times. 4000 g — 20 min.
3. Store it in refrigerator for overnight at ca. 5°C. (Ca. 20 hrs)
4. Treat with duponol (30 ml : 15 % : at room temp :) for 5 hrs. pH. 7.5
   7:10 ⟶ 12:10
5. p.p.t with 95 % of Et-OH.
6. Wash the p.p.t with 80 % Et-OH. twice.
7. Suspend the p.p.t in saline-citrate soln with aid of glass-homogenizer 60 ml.
8. C.f.g. 3,000 xg : 20 min and repeat this twice.
9. Sediment saved.
10. Extracted twice in 40 ml of 2 M NaCl on Magi Mix. at 4°C for 30'.
11. C.f.g 20,000 xg for 20 min. at 4°C.

23/III

12. Add it to 100 ml of 95 % Et-OH.

24/III

13. Fibers suspended in 5 ml of 2 M NaCl.
14. Deproteinize with sevag's method : Add equal volume of CHCl$_3$ + Amyl alc. mix. Shake it for 5 min. Repeat [redacted] times. No interface.
15. P.p.t with Et-OH [redacted]
16. Dissolve in small 2 M NaCl. soln.
17. Digest RNA with RNase. (50 γ/ml. final. : 1 hr at 37°C.)

1 mg/ml Pronase 10 min : 37°C

U.W.
MICROBIAL
GENETICS

Mating capacity of depronolated 3747.
(15% dupond).

11/III, 1960 REF:

Method :

Use 20 ml culture of 3747 (phenary grown for overnight at 37°C)
(2 tubes of 10 ml)
Recipient 4573.

↓ C.f.g.

↓ resuspend it in 1.5 ml of H_2O.

↓ Add 1 ml of the suspension to 5 ml of 15% dupond.

↓ keep it at 37°C.

Take 1.0 ml of the sample, add 2.0 ml of H_2O
↓ C.f.g. {12000 rpm at each time 0 hr, 1 hr, 3 hr.
20'
wash it once with 5 ml of H_2O.

↓ resuspend it in 1 ml of H_2O.

Use this as a depronolated cell.

↓ Use 0.2 ml.

↓ Test mating capacity. for survival, for mating, for DNase digestion.

0.2 ml 3747 depronolated + 4573, 1 ml + Pan 8 ml.
Exp. grown.

Result :—

Volume of depronolated 3747 susp. used	3:00	4:00 +DNase		5:00	No DNase added.		
	1 hr	2 hr	3 hr		1 hr	2 hr	3 hr
# of survival colonies on EMB Lac 10^{-7} / 0.2 ml. of the treated suspension.	17, 20	4, 2	0, 1		14, 12	5, 2	4, 3
# of Recombinants / plate. 0.2 ml / plate. on Mlac Str.	0, 0	0, 0	0, 0		0, 01	0, 0	0, 1

U.W.
MICROBIAL
GENETICS

Mating capacity of duponolated 3747. (10% duponol).

12/III ; 1960

REF:

	1	2	3	4	5	6	7	8	9	10
		• Cultural age :		overnight culture of 3747 in Penassay broth. (30 ml)						
		Recipient :		4573.						
		• Treatment :		10% duponol : final conc. at 37°C.						
		• Time :		1 hr. (2:30)	2 hr. (3:30)	3 hr. (4:30)				
		• c.f.g. :		aerval ; 12000 rpm. for 20 min.						
				3.5 ml H_2O + 1.5 ml. duponolated 3747						
				Wash it with ↓ 5 ml of water						
				Resuspend it into ↓ 1.5 ml of H_2O.						
		• Result :								

U.W.
MICROBIAL
GENETICS

Mating capacity of duponolated 3747.
(10% duponol)

12/XII ; 1960

REF:

1	2	3	4	5	6	7	8	9	10
	• Cultural age :		Overnight culture of 3747 in Penassay broth. (30 ml)						
	Recipient :		4573.		"				
	• Treatment :		10% duponol : final conc.		at 37°C.				
	• Time :		1 hr. (2:30)	2 hr. (3:30)	3 hr. (4:30)				
	• c.f.g. :		serial ; 12000 rpm.		for 20 min.				
			3.5 ml H_2O + 1.5 ml duponolated 3747						
			Wash it with ↓		5 ml of water				
			Resuspend it in ↓		1.5 ml of H_2O.				
	• Result :								

U.W.
MICROBIAL
GENETICS

Transformation in E. coli K-12.
using Helper (Hfr, F+, F−)

14/III ; 1960 REF:

• Strains used:

DNA from 3747 : 1 month old. (Feb. 10. in 2M saline. kept in refrigerator.
Recipient ; 4573 : F− Lac85
Helper : { 4574 F+ : F+ Lac85
4574 : F− Lac85
4049 ; Hfr, (M P V_0^R Lac$_{89}$ Hfr,) m.

(Those culture was taken from old broth culture (several weeks)).

• Cultural condition:
Inoculum size : 0.2 ml / 8 ml penassay. ; Incubate at 37°c on rotator. 11:00 ~ 2:00
Use exponentially growing cells.

• Experimental condition:

Inoculate 0.1 ml of Recipient
+
0.1 ml of Helper
+
0.1 ml of DNA. (ca. 20γ /ml)
+
1.5 ~~1.8~~ ml of Penassay broth.
(make whol volume into 2 ml.). 2:15 ~ ca. 10^9 ~ 10^{10}.

→ Incubate it at 37°c for overnight on rotator. → Seed it on Hfac Sm.

Use 0.2 ml / plate EMfac Sm.

• Result : After overnight shaking concentration of cells become

Exp. #	Mix: Recipient	Helper	DNA	# of lac+ x+ on Hfac Sm
1.	4573 +	4574 +	3747	0, 0
2.	4573 +	4574 + 4574F+	3747	0, 0
3.	4573 +	4574F+	3747	0, 0
4.	4573 +	4574F− 4049	3747	0, 0
5.	4573 +	4049	3747	0, 0
6.	4573 +	4574 4574F+ 4049	3747	0, 0
7.	4573 +	4574F+ 4049	3747	0, 0
8.	4573	———	3747	0, 0

Test sterility of DNA. on Hfac. 0.1 ml / per tube : OK ; Steril after overnight shaking on rotator which is same as 1 ~ 8.

order of additions of the DNA cultures: I II III.

U.W.
MICROBIAL
GENETICS

DNase-sensitivity of λ mating capacity of Sm-killed cell. a

14/III ; 1960 REF:

Experiment : Strain. : 3747 overnight culture at 37°C.
↓
+ Sm 1 mg : final conc.
↓
keep ~~5~~ incubation overnight.
↓
c.f.g & resuspend it in 5 ml of Penassay broth.
↓
0.2 ml bacterial suspension. + 0.1 ml enzyme soln.
+ DNase 200 γ/ml
+ RNase 100 γ/ml
+ lipase.
4:00 ~ 5:00.
Incubate it at 37°C for 1 hr.
↓
+ 4573 1 ml + Penassay 1 ml.
↓
Incubate the mix for 1 hr. at 37°c.
5:00 ~ 6:00
↓
plate the mix onto MLacSm. 10^{-1} : 0.1 ml / plate.

Result :

	● No enz.	+ DNase	+ RNase	+ lipase
# of Recombinants / plate EMLacSm 10^{-1} : 0.1 ml				
# of survival cells/ml on EMB lac. 10^{-1} : 0.1.				

Conclusion : seems no difference.

U.W.
MICROBIAL
GENETICS

Transformation of E. coli K-12.
(Hfr, F+, F- Helper). without shaking.

15/III. 1960 REF:

Lac$^{-}_{85}$ → Lac^{+}

· without shaking. Idea : DNA may be taking up at the time of infection of F+ from F+ to F-

· Cultural age of strains used.

10:30 ~ 1:30
0.2ml / 10ml pen. → at 37°C.

· conc. of DNA. 200γ/ml : final concentration.

· Experimental condition for transformation of E. coli.

Mix these bacteria and ONA.

0.1 ml	4049 Hfr,	} helper.	final Ca. 10^8 cells/ml for each strain.
0.1 ml	4574 F-		
0.1 ml	4574 F+		
0.1 ml.	4573	} : Recipient	
200γ/ml : 0.2 ml/2ml		: DNA.	

(kept in refrigerator for 1 month)

Make this mix into 2ml with pen. broth.

↓

· keep incubation for 3.0 hrs / overnight at 37°c without shaking. (just stand the mix.)

↓

· Score Lac^{+} S^{R} X^{+}.

Inoculate 0.2 ml of these cultures on M Lac Sm agar.

· Result :

Vol. of Pen. (ml)	Exp. #	Constitution of the mix: Helper	DNA	Recipient	# of transformants/plate 3.0 hr	24 hr
1.6	1	4574 F+	3747	4573	0, 0	0, 0
1.5	2	4574 F-	3747	4573	0, 0	0, 0
1.5	3	4049	3747	4573	0, 0	0, 0
1.4	4	4574F+ + 4574F-	3747	4573	0, 0	0, 0
1.4	5	4574F+ + 4049	3747	4573	0, 0	0, 0
1.4	6	4574F- + 4049	3747	4573	0, 0	0, 0
1.3	7	4574 F+ + 4574F- + 4049.	3747	4573	0, 0	0, 0
1.6	8	——	3747	4573	0, 0	0, 0
1.4	9	4574F+ + 4574F- + 4049	—	4573	0, 0	0, 0
10 Lac		3	1	2		

Conclusion : No transformants were observed under this condition.

U.W.
MICROBIAL
GENETICS

Extract DNA from W4687.

(Use this DNA as Gal-transfer.)

18/III. 1960 REF:

EML.

1. Grow up cells in penassay broth (Inoculated from broth given by Esthy)
10 ml. : overnight grown.

2. Inoculate the 10 ml into 2 l of Penassay in 6 l flask.
Shake it for overnight at 37°C. 2:00 ~

3. Collect cells by centrifugation.

· Continuous flow.
refrigerating centrifuge.

· Bactogar.

U.W.
MICROBIAL
GENETICS

Isolation of DNA from 4687.

$P_1 \rightarrow$ × 4320
→ × 4621 $F^- Ind\ 4^+ bl_{27}^{--} S^R$
#74. (bal +/− P_1^+? 9 4^+? F^R

19/III :1960 REF:

1. Strain W4687 grown in pen at 37°C with heavy shaking. (2
   for 6 hrs. on shaker. (2.5 l. in 6 l flask
2. Collect cells with Sorvall centrifuge : 4000 rpm. for 10 min
   wash the cells twice with 0.14 M saline. Ca. 3 ~ 4 grs ?
3. Let it suspend in H_2O by addition of 2 ml H_2O.
4. Keep it in refrigerator : 9:30 PM ~ 7:30 AM.
5. Add the bacterial suspension to 30 ml of detergent soln (15% pH 8.8)
6. Keep it at room temp for 5 hrs (7:30 ~ 12:30).
7. p.p.t. with 95% Et-OH. (4°C)
8. Wash the p.p.t. with 80% Et-OH. twice.
9. Suspend the p.p.t in Saline-citrate soln. with the aid of glass homogenizer (Ten Broeck)
   (0.15 M) (0.005 M)
10. C.f.g. & save the ~~pellet~~.
11. Extract DNA with 40 ml of 2 M NaCl (twice) on Magi Mix. at 4°C for 30'
12. C.f.g at 20,000 xg for 20' at 4°C.
13. Add the supernate to 100 ml cold 95% EtOH.
14. Suspend p.p.t in 7 ml of 2 M saline.
15. Keep the solution at 5°C for overnight.
16. Deproteinize with $CHCl_3$-octanol method (5) : (1) { $CHCl_3$ + octanol / = DNA soln /
    Make 5 x Seveg.
17. ~~Deproteinize~~ p.p.t : DNA with 80% Et-OH.
18. Dissolve it in 5 ml 2 M NaCl. → Sample I.
19. Take 2.5 ml of the ~~solution~~ solution (twice) (Sterilized) and add 50 γ/ml of RNase (fresh Conc.)
    Keep at 37°C for 60'. (Boiled)
20. p.p.t. DNA with 80% Et-OH and dissolve it in 2 M salt. → Sample II.

11/18/59 EM259.67

P-1 + W4520 $\mathcal{N}_{Fe}$

——× W4621
$F^- Ind^- Lp^+ Gal_2^- y^- S^R$

transductants 73 & 74

Gal +/−
F'8
P_1 + ?
Lp + ?
S^R Ind-

recorded 3/15/60 as W4687
#74
as Gal +/- heterogenote.
(Segregation Pattern: details later)

♀ $Gal_6 S^S$ 3096
S^R 3996

U.W.
MICROBIAL
GENETICS

DNA-analysis.

20/III ; 1960 REF:

Ref: Burton, K. : Biochem. J., 62, 315~323 (1956)

Reagent:

1. Diphenylamine Reagent.

Dissolve 1.5 gr of Fisher certified diphenylamine in 100 ml of redistilled acetic acid and add 1.5 ml of conc. H_2SO_4. Store in brown bottle.

On the day it is to be used 0.10 ml of aqueous acetaldehyde (16 mg/ml) is added for each 20 ml of reagent req'd.

Procedure:

The author states that extraction with 0.5 N $HClO_4$ at 70° for 15 min. (2x) will liberate 95% of the nucleic acid from E. coli. This is comparable to extracting at 90° for 15 min. with 5% TCA.

The final conc of $HClO_4$ must be 0.5 N before add'n of diphenylamine reagent.

1.0 ~ 2.0 (usually 1.5 ml) of sample is mixed with 2 volumes (3 ml) of diphenylamine reagent containing acetaldehyde.

Tubes containing known amounts of Standard DNA and a blank containing 0.5 N $HClO_4$ but no DNA are also prepared.

Color is developed by incubating at 25~30°C for 16~20 hr. Provided that all tubes are at the same temp, constancy of the temp. is not essential, and variations at least between 25°C and 35°C do not appreciably affect the readings.

Read at 560 mμ.

Added. read at ~ P.M.

Standard: 5γ, 500γ/ml of DNA.

Use 0.1 ml (5N) P.C.A to 1.5 ml of sample and 3 ml

Volume of Sample used.

conc. Standard		ml Sample I		ml. Sample II	
50γ	5γ	0.1	0.3	0.2	0.3
.85	.075	0.36	1.9	0.24	2.0

0.1 20 ml. EN 0.045
1.5 ml
0.015 3 ml.

Exp.:-

0.1 ml + 1.4 ml H_2O + 0.15 ml PCA (5N)

0.3 ml + 1.2 ml H_2O + 0.15 ml P.C.A (5N)

↓

+ 3 ml diphenylamine reagent cont. CH_3CHO.

Standard (desoxyadenylic acid. 5γ, 500γ/ml.)

0.1 ml + 1.4 ml H_2O + 0.15 ml P.C.A (5N)

blank — + 1.5 ml H_2O + 0.15 ml PCA (5N)

U.W.
MICROBIAL
GENETICS

Pentose determination.

29/III ; 1960 REF:

Ref. Colowick & Kaplan "method in Enzymology" Vol III, p. 87.
(Dische modification: J. B. C., 204, 983 (1953))

Reagents:

Acid Reagent – 10% $FeCl_3 \cdot 6H_2O$
0.5 ml of $FeCl_3$ solution in to 100 ml of Conc. Hcl.

Orcinole Reagent – 6% soln of orcinol in 95% Et. OH.

Procedure:

To 1.5 ml of unknown is added 3 ml of the $FeCl_3 \cdot 6H_2O$ (acid reagent above).

Add 0.2 ml of freshly prepared Orcinol soln

Heat in Boiling H_2O bath for 20 min.

Read in Klett with #66 filter.

Comments:

Under these condition, both ribose & ribose-5-PO4 are completely developed in 20 min.

No other sugars besides pentoses can be present.

Standard soln of ribose used as control.

Standard: 5 to 50 γ/ml : final.
(1/10)
ribose. soln is used as standard

Vol. Sample (ml)
0.1 ml

U.W.
MICROBIAL
GENETICS

2nd Extraction of DNA From 4687.

25/III ; 1960 REF:

1. Grow up cells in penassay (2.5ℓ) on rotary-shaker for overnight at 37°C. ~1ℓ flask.
2. C.f.g. & wash two times. with 0.14 M saline. Store in refrigerator at 5°C for overnight. 25/III ↑
3. Add 1% (pH. 7.5 ajusted 0.5ml of 0.01 N HCl) @, 30 ml of Onpond sol. and keep it at room temp for 5 hrs. 7:30 ~ 12:30
4. p.p.t with 95% of Et-OH.
5. Wash the ppt with 80% Et-OH twice.
6. Suspend the ppt in 60 ml of saline-citrate sol. with aid of glass-homogenizer.
7. C.f.g. (30,000 xg : 20 min) Sediment saved. Repeat this three times.
8. Extract twice in 40 ml of 2M NaCl on Magi mix at 4°C for 30'.
9. C.f.g (20,000 xg : 20 min) at 4°C.
10. Add it to 100 ml of 95% Et-OH. 26/III ↓ 80% 200 ml = 95% 168 ml + H_2O 32 ml
11. Fibers suspended in 10 ml of 2M NaCl.
12. Deproteinize with Sevag's procedure: $CHCl_3$ 5 : N-octyl alcohol 1 Mix 1 : 1. Repeat 5 times.
    After 7& times Sevag, there is no interface.
13. p.p.t with 100 ml, 95% Et-OH.
14. Dissolve in 10 ml of 2M saline. ——4687 I

U.W.
MICROBIAL
GENETICS

Extraction of DNA from 4520. ($H\ \lambda_L^2\ F_8$)

21/I ; 1960 REF:

1. Grow up cells for 6 hrs in 2.5 l pan (in 6 l flask) with shaking at 37°C.
2. Wash cells 3 times with 0.1 M saline. keep it in refrig. at 5°C for overnight
3. Treat with 15% duponol at room temp for 5 hr. 7:30 AM ~ 12:30
   cell suspension was not well homogenized : looks bad.
4. p.p.t. with 95% Et-OH.
5. Wash ppt with 80% Et-OH. twice
6. Suspend p.p.t in Saline-citrate soln. with the aids of glass-homogenizer
7. C.f.g. it (@ 30,000 xg). Saved p.p.t. Repeat 2 times.
8. Extract DNA with 40 ml of 2M saline. (2 times.) (20 min extraction at 5°C with Mag. Stir
9. p.p.t. DNA with 95% Et.OH. 100 ml.
10. Dissolve the p.p.t. in 2 M salt. Store in refrigerated at 5°C.
11. Deproteinize it 4 times by Sevag's procedure. there is very little interface insoluble remain.

    In the course of this procedure, the sample was kept at room temp. For 1 hr.
12. p.p.t with 100 ml of 95% Et.OH.
13. Dissolve it in 5 ml of 2M saline. (steril.)
    • this sample seems more easily dissolved in 2 M salt than the other samples. Also seems very thick.

U.W.
MICROBIAL
GENETICS

Biological activity of DNA extracted from several Strains. (Gal⁻ → Gal⁺ : plaque forming activity).

1/11 ; 1960 REF:

1	2 Lysogenic for	3	4	5	6	7 Condition:	8	9	10
Recipient :	434 hy.	: Gal+		Helper : 434 hy.		at 30°C. for 1 hr.			
plate #	Strain Extracted DNA	0.1 ml DNA 0.1 ml Bact.	0.1 ml DNA : 1 ml of Media		# of plaques	# of gal+ on T.T.C & on MGal (T.T.C : soft agar)			
1	4687 "New"		0.1	TTC	0	37 ?			
2			0.1	EMGal	0	0			
3				TTC	Ca. 1000	0			
4				EMGal	527	0			
5	from 3094.		0.1	TTC	4	1			
6			0.1	EMGal	1	2			
7	$4^+ 4/4r^2$ (dg)			TTC	480	0			
8				EMgal	199	0			
9	4520		0.1	TTC	4	1			
10	4^+ Gal$^+$ / Gal$^+$ F_8		0.1	EMGal	10	0			
11	(Through this DNA seems no good must be repeated.			TTC	3	4			
12				MGal	0	0			
13	old 4687		0.1	TTC	3	6			
14			0.1	EMgal	5	1			
15				TTC.	395	2			
16				Mgal.	46	0			

Conclusions:

1. DNA from 4687, & 3094 have plaque-forming activity.
   a. It forms plaques.
   b. Such activity will disappeared by DNase-treatment.
   c. That activity is retained or in preparation during the process of preparation of DNA
2. There is no transforming activity for Gal.

U.W.
MICROBIAL
GENETICS

2nd Extraction of DNA from 4520.

2/14 1960 REF:

1. Let 4520 grow for overnight on rotator.
2. Collect cell with 0.14M saline. Wash 3 times with saline.
3. keep it at 5°C for overnight.
4. lyse cells with Duponol (15%) for 5 hrs. at room temp.
5. ppt. & rinse it with 95% Et. OH
6. Wash cells with Saline-citrate mix 60 ml. 3 times.
   0.14M 0.05M.
   Suspend the ppt in Saline-citrate mix. Mix with Mag-mix. for 20 min.
   c.f.g. x 50,000 g for 20'. Save sediment.
7. Extract DNA with 2M saline (40 ml : 2 times) with Mag.mix. twice
   30'. at 4°C. 2:15 ~ 2:45.
8. c.f.g. 20,000 xg for 20'. Sup. saved.
9. Add. it to 100 ml of 95% Et-OH.
10. Dissolve it in 10 ml of 2M NaCl.



U.W.
MICROBIAL
GENETICS

W 3104 F_8

11/14. 1960 Made by 4520 F_p × 3104 REF:

Test of compatibility.

4520 × 3104



3104

4520

m Mal+

Gal+ F_8

1. W3104 F_8 : grow up cells in 10 ml broth. 4. Add it in ~~25l~~ pm. Shake it for 5 hr. on rotator.
2. C.f.g & collect cells by small c.f.g. wash it twice with 0.14 M saline.
3. Treat with 15% dupanol at room temp for 5 hrs. 10:30 AM ~~—~~ 3:30 pm.
4. P.P.T. DNA with 95% Et-OH (100 ml) Rinse it twice
5. Wash p.p.t. with citrate - Nacl-Soln. twice, Save sediment. Homogenized by potter-type homogen.
6. Extract DNA with 40 ml of 2M Nacl twice.
7. P.P.T it with 100 ml of cold Et-OH. Dissolve it in 10 ml of 2M Nacl

U.W.
MICROBIAL
GENETICS

B 114 (W3104 1dg) lyse by U.V. No λ release
Immune to λ infection.

15/11 1960 REF:

1. Grow up cells in 2.5 l pen. for overnight at 37°C.
2. Collect cells keep it in refrigerator for overnight. (5°C).
3. Lyse the cells with 15% deoxycholate at room temp (11:00 ~ 4:00) for 5 hrs.
assay of plaque forming activity. 10^{-1} : 0.1 / plate on 3110.
4. Wash the cells with ~~95%~~ 70% Et-OH. (100 ml.) $CHCl_3$-treated. — c.f. g. - Store in refrig.
5. Wash the precipitate with citrate-saline soln. twice.
6. Extract DNA with 2M saline : 40 ml : twice. on Mag. mix. for 30'. at 5°C.
7. P.P.T the extract with 95% Et-OH. (100 ml), & Dissolve it in 2M saline (10 ml)
8. Deproteinize it with Octanol-$CHCl_3$ soln. (5 times) until no interface
9. P.P.T DNA with Et-OH and redissolve it in 5 ml of 2M saline.
10. Store in refrigerator at 5°C.

Test for λ production.



No plaque was found on 3110.

U.W.
MICROBIAL
GENETICS

Labeling DNA with P^{32}

1960. REF:

Medium low in phosphate for making P^{32} DNA in E. coli.

Ref. Lehman, Doesman et. al. : J. B. C., 233, 163 (1958)

The medium contains .34 μM of $PO_4^{\equiv}$ / ml.
Coli requires ca. .70 μM/ml for optimal growth.

Preparation of Medium.

Na-lactate 70 mMol.
to 100 ml of hot lactic acid (Mark 85%), add 40% NaOH (~100 ml) to pH. 7. Boil 3 min. Check pH.
Cool & filter if necessary. Soln is yellow viscus.
concentration ~ 6 M (~100 ml)

glycerol	4g
NaCl	5 gr.
KCl	2 gr
NH_4Cl	1
$MgCl_2$	1 mMol
$CaCl_2$	0.1 μMol
gelatin	0.01 gr
P (as ortho PO_4)	10 mg P
S (as $SO_4^{\equiv}$	10 mg S

To 1,000 ml should be pH. 7.
generation time : ~ 1 1/2 hr.
cells smaller than in most of the synthetic med.

Preparation of Stock Soln.

A.) Na-lactate ~6 M 58.0 ml
glycerol 95% 16.0 ml
NH_4Cl 5.0 g
→ 200 ml pH. 7.2

B)

NaCl	25g		
KCl	10		
$MgCl_2$	0.3 M	16 ml	
$CaCl_2$	1 M	0.5 ml	
gelatin	1%	5 ml	
KH_2PO_4	pH. 7.4	1 M	1.7 ml

to (5 ml) 200 ml. pH. 7.3.

C). Na_2SO_4 - 0.2% After autoclaving may be mixed with B (after B has been autoclaved)

Autoclave all 3 stock soln 20'

Preparation of complete medium.

92 ml sterile H_2O (+ P^{32})
4 ml A
4 ml B
4 ml C.

● continued to back page

prep. of P^{32} med. — Want 0.6 ~ 0.8 μM P/ml

Add hot P^{32} + adequate cold KPO_4 1 M.

pH. ~7. to ~~adjust~~ give difference from 0.34 μM/ml for E. coli

for P^{32} studies, the P^{32} should be added to the water, autoclaved, and then neutralized with sterile NaOH.

(Best autoclave ~~with~~ while P^{32} in dilute acid, since any polyphosphates will then be broken down.)

Inoculum — can be grown in Bunsen. Use a small inoculum, and grow with vigorous aeration.

U.W.
MICROBIAL
GENETICS

Labelling cells with P^{32}.

c.f. Ferro, F

13/IX ; 1960 Exp. cell. Res. 12, 363-373 (1957) P^{32} : PO_4 in weak HCl conc. REF: ~~Specific activity~~ : 200 mc/ml.
Specific activity : 55350 mc/g.
date : 8-29-60.

purpose : 1. for autoradiography of bacteria.
2. Testing leakage of P^{32} from P^{32} labelling bacteria after dilution.

Method ① grow up cells in H. medium for overnight at 37°C on rotator.

② ~~Wash the cell~~ Add the 10ml of the culture to 3 med. made by method as follows.

a. To 0.8 ml H_2O cont. (1/40 conc ord.) add 0.01 ml (10%) stock.
2% CH : 0.1 ml / 10 ml H med. : final conc. : 0.2%.
0.5% CH : 0.1 ml / 2 ml H : 0.25%

b Evaporate it and redissolve it in 0.8 ml H-medium
adjust pH. 7.2. Add. 0.1N KOH ca. 0.05 ml.

③ Grow cells for 3 hr on shaker at 37°C.
1:100 ~ 1:500 Inoculum size Ca. 10^{7}
10% : .8 ml.

④ Wash cells 5 times. with ~~cold~~ water : 2 ml.

No. of Plate ①	# of Washing	before wash (total) counts/min. ×1000	
2	0	(Sup.)	50% 120,000
3	1		×10 1200,000
4	2	4553	
5	3	1493	10^{-3}
6	6 (suspended in 2 ml)	12217 ×5 = 61085 count/min	

⑤ ~~Add~~ Add 0.25 ml of the suspension to 10 ml of protoplastic broth (c.f. back page) shake it on rotator.

⑥ Make high speed c.f.g. Save sediment. Add 1 ml of H_2O to both tube. Assay Radioactivity in sediment.
50,000 ×g : 20'
(Keep it 15° after + H_2O) c.f.g ×3,000×g

	7 + P	8 P (×1000)
Count/.1 ml/min.	2284	2248

1200

1