Friday 10/10

- Vg/6 with A_2 - that failed to grow on individual A_2's both added to result of streaked out CHBloc -

- 811 t (750t1821) ~~Transduced~~ 8 papillae picked and streaked for stability check
  IS TRANSDUCTION PRODUCED BY TRANSDUCED STRAIN STABLE? y — — — y 129

- 578 t (750t1821) 11 as 811 above.

- 811 gut R- 2 lysate prepared
  filtered
  sterility checked (0.9 ml) in broth - no growth

Sunday. 10/12

S10 absorption Expt- plated on EMB bac-, all others bac as expected

(1.) cell stock assay = $X10^{8}$, — 67,74 = 7.1×10^{8} cells/ml

(2.) phage stock assay = $X10^{8}$ — 531,453 = 4.9×10^{9} phage part/ml
[a 2+ hour broth del. of phage stock IC-12 25 λ]

INPUT RATIO PHAGE/BACTERIUM $\frac{49}{7.1} = 7/1$

(3.) phage stock gal'dent/0.1 ml = 177, 176 = 1.77×10^{3}/ml - spontaneous

(4.) Spontaneous gal+/0.1 ml cells plated = - 15

1.62×10^{3} gal'dent/ml λ susp.

gal'd/ml × dil = original titer

$1.62^{+3} \times 6 = 7.72 \times 10^{3}$/ml original stock

previous assay 125 = $6.91^{+} \times 10^{3}$
higher titer likely probably due to increased accuracy of longer dil. of lysate

Gal'd -

Ratio gal'd $\frac{9.72 \times 10^{3}}{4.9 \times 10^{9}} = \frac{2}{10^{6}}$

$= 1/5 \times 10^{5}$ λ

adsorpt = 5 min bench
10 min centrifuge

2 ml cell susp + 2 ml dil. λ

(5.) Supernat. 1st adsorption λ titer = $X10^{7}$ — 72,126 = 9.9×10^{8}

added = stock titer / 2 = $\frac{4.9 \times 10^{9}}{2} = 2.5 \times 10^{9} = 25 \times 10^{8}$

unabsorbed = 9.9×10^{8} = 39%

λ taken up/cell = $\frac{25 \times 10^{8} - 9.9 \times 10^{8}}{3.6 \times 10^{8}} = \boxed{4.1 \text{ λ/cell}}$

1st

(6.) Sediment 1st adsorption - no gal'dents

0.1 ml plated = 124, 138 . 131

gal'ded = 131 - bkgrd. = 131 - 15 = 116

expected from above stock assay = 81 ← indicates more than 100% adsorpt

(7.) supernat 1st adsorpt.

0.1 ml plated = 33, 33 = 33

gal'd. present = total pop. - sp = 33 - 15 = 17

(8.) supernat 2nd adsorpt λ titer = $X11^{7}$ — 144,103 = 1.24×10^{9}

put in = 2.5×10^{8}
unabsorbed = 1.2×10^{9} = 50%

absorbed/cell = $\frac{25-12}{3.6} = \frac{13}{3.6} = \boxed{3.6 \text{ λ/cell}}$

total = 7.7 λ/cell

1st adsorption - amount of killing by λ

cell assay = $X10^{7}$ = 139,132 = 1.56×10^{9} cells/ml
which appears impossible

continued

Sunday 10/12

(9) 2nd adsorption ~~Sed~~ sup. no of gold.

0.1 ml plated = 169 149 = 159

gold included = total - sp = 159 - 15 = 145

expected max if 100% adsorpt.
on basis initial assay = $\frac{16^2}{2} + \frac{16^2}{2} = 162$

expected on basis of adsorpt. of 1st & 2nd = 81 (.61) + 81 (0.56) = c. 90

(10) 2nd adsorption supernat. no of gold.

0.1 ml = 62, 64 = 63 - 15 = 48

gold. added /ml = 81 81

gold. supernat = 63 48 } total 92. ↑↓ comp.

gold sed = ~~145~~ - 116 = 29

(11) 3rd adsorpt. ↓ titer supernat. = $\times 10^7$ = 226, 198 = 2.1×10^9

input = 2.5×10^9 /ml

adsorbed = 2.1×10^9 /ml = 84%

adsorbed $\frac{0.4 \times 10^9}{3.6 \times 10^8}$ = 1.1 /cell

λ/cell

↑

This value may be off due to the time elapse between the beginning of this expt and this assay - Time is probably in excess of 1 hr and therefore possibly longer than latent period.

total = 7.7
1.1
8.8 λ /ml

(12) 3rd adsorpt. sed. no of gold.

0.1 ml = 179, 214 = 197

increase over 2nd 197 - 145 = 52.

(13) 3rd adsorpt. sup. no of gold.

0.1 ml = 53, 65

gold = total - sp = 59 - 15 = 44

total gold = 52
44
96

expted (81)

81
44
37

Thursday 10/16/52

- 518 Second absorption Exp. -

518 blkgnd was ~ orig. plaque assay 41

- Initial phage stock titer - a 1-2 dil. of parent 25'
  10^8 dil → 80,74 = 7.7×10^9 λ/ml (parent assay 1.6×10^{10})
- Initial cell assay - stock susp. from overnight cult. resusp. in saline
  10^7 dil → 115,108 = 1.1×10^9 cells/ml

Expt. performed by adding 2ml volumes of λ + cells together 5 min at rm temp. Centr. 10 min. Decant sup, resusp. cells in 2ml volume - Remove samples, i.e., 0.2, 0.1 ml - making vol. either 1.8 or 1.9 ml - For next absorpt add 1.8 or 1.9 ml of λ sup., etc.

- Initial phage stock gold titer
  0.1 ml → 457 pfu - 41 sp = 416

1st absorpt.

Supernat λ titer -
10^7 dil → 164,205 = 186×10^7
% not absorbed = $\frac{3.9 \times 10^9}{7.7 \times 10^9}$ = 48% absorbed = 52%

Cell survival (of 2nd resuspension)
10^7 dil 134,89 = 112×10^7 = 1.1×10^9 cells = no killing

Supernat gold.
0.1 ml = 197,166 = 182 − 41 = 141 % gold not absorbed: $\frac{141 \times 2}{416} = \frac{282}{416}$ = 67.5%

Sed. gold.
0.1 ml = 268,317 = 293 − 41 = 252

Vol. at end = 4.0 ml − 0.1 (sup) − 0.1 (sup of gold) − 0.1 (sed gold) = 3.7

2nd absorpt

Supernat λ titer.
10^7 dil → 232,329 = 2.8×10^9
% not abs. = $\frac{5.6 \times 10^9}{7.7 \times 10^9}$ = 73 = absorbed = 27%

Supernat gold.
0.1 ml = 309,281 = 295 − 41 = 254

Sed gold.
$\frac{0.1}{5}$ = 107,79 = 93 − 41 = 52

Vol. at end: 3.7 − 0.1 − 0.1 = 3.5 ml

3rd absorpt

Supernat λ titer
10^7 dil → 215,342 = 2.79×10^9
% not abs. $\frac{5.6 \times 10^9}{7.7 \times 10^9}$ = 73 = absorbed = 27%

Supernat gold.
0.1 ml = 368,345 = 347 − 41 = 306

Sed. gold
$\frac{0.1}{5}$ = 129,85 = 107 − 41 = 66 × 5 = 330

4th absorpt

Supernat λ titer
10^7 dil → 253,293 = 2.7×10^9
% not absorbed = $\frac{5.4 \times 10^9}{7.7 \times 10^9}$ = 70 absorbed = 30%

Supernat gold
0.1 = 283,246 = 242

Sed. gold
$\frac{0.1}{5}$ = 43.49

Thursday 10/11/52

5th adsorpt

Supernat titer

10^7 dil → 552, 527 = 5.4×10^9 = ~~4.2X10⁸~~

% not adsorbed = $\frac{1.1 \times 10^{10}}{7.7 \times 10^9}$ = impossible - either no adsorpt or lysis obscuring effect.

Supernat gold.

Sed. gold.

Sed pt. assay

10^7 dil → 44, 51 = 9.6×10^8

~~~~

- 1924 Examination for presence of λ

14 cross streaks of overnight per culture streaked against 1985

9 showed presence of phage - weak.

- 811 - gel + - 2 λ neg.

10^5 dil = 29, 23 = 2.6×10^9 λ/ml

- 1103 - Gel

		15	← 1.6×10^{11} λ/ml
		90	←

- 513 Gel

0	811 - gel +	27
1	811 gel +	1
5	811 λ	171
	811 λ -	29

(broad)

- 2000
~~~~

Friday 1/11/[illegible]

- 515 (Lgp.sta.o. [illegible]) X 12 - [illegible]
  [illegible] gel +
  [illegible] [illegible] - [illegible] / [illegible] - [illegible]

- 519 + ([illegible]) [illegible]
  [illegible] s/g mixed
  [illegible] s/g [illegible]

- 811 ([illegible])
  [illegible] s/g mixed
  [illegible] s/g mixed

- W1655 X W702 - Lgp[5] gel - transmissible gel
  tested on mixed - is mixed +

- Research on Sycogenicity of 1924
  - 12 cross streaks on 1485 - no evidence of phage
    on gal, gal -
    on EMB lac, lac -

- 1924.
- Examination of 1924 for lysogenicity of transinduced cells. -

Isol. #	Streaking # ①	②	③	Streaked against 1485
1	m	m	m ⊗	non-lyp.
2	m	+	+	" "
3	+	+	+	" "
4	m	+	+	"
5	⊗ m	m	m ⊗	one plaque
6	⊗ m	+	+	non lyp.
7	m	+	+	"
8	m	+	+	"
9	+	+	+	"
10	m	+	+	"
11	+	+	+	"
12	m	M	m	"
13	+	+	+	"
14	+	+	+	"
15	+	+	m	"

Isol. #	①	②	③	/1485
16	m	+	+	non lyp.
17	m	+	+	"
18	+	+	+	"
19	m	+	+	"
20	m	m	+ ?	"
21	+	+	+	"
22	+	+	+	"
23	m	+	+	"
24	m	+	+	"
25	m	m	⊗ m	"
26	m	+	+ ?	"
27	m	m	m	"
28	m	+	m	"
29	m	m	m	"
30	m	m	m	"

Isol. #	①	②	③	/1485
31	m	+	+	non lyp.
32	m	m	m	"
33	+	+	+	"
34	m	+	+	"
35	+	+	+	-
36	+	+	+	"

μ = mixed
+ = gel
- = -
⊗ = mosaic

Sunday Oct. 19

- 1821 - Reexamination of xylose transduction possibilities

EMB gal	Additions	no gal 2 days	
1	—	174 +	8 picked and streaked -
2	0.1 K gal (25')	229 +	16 picked and streaked -

Examination for instability of "gal deficient" -

- Plan to gal deficient 1821 and examine unstable gal+ for gal stability -

- 1924 H 1242 see p. 129

- 2050 gal$_3$ - mutants - streaked on EMB base for replication on mucinol + supplements

- out of 12 suspected auxotrophs - 1. prototroph
  1. A_2
  1. ($A_1 A_2 A_3$) - really proto! failed on A_5 } not tested A_4 - presumably same as A_2

SIP

- Check on background on EMB gal.
  each from overnight culture - uncorrected

EMB gal	Additions	no gal 2 days
1	mucinol. supp	53
2	1-10 supp	37
3	1-100 supp	43

(probably)

811-gal+

- Crosses - X 1295 - Results partly unsat. due to large no. of mucoid col. - many mucoid apparently (-)

811-gal+ 2 - (no galactokin in 811)

EMB gal	+	supp -	checked -
1	33	3	
2	39	2	
3	29	0	
4	46	0	
5	34	0	
6	37	0	
7	48	1	
8	23	0	
	289	6	6/6 gal slow

811 gal+ 1

1	956	1
2	71	2
3	70	0
4	62	0
5	67	0
6	64	3
7	66	2
8	78	1
9	84	2
10	66	1

- Wednesday -

1821 - Xyldirection? —

	EMB xyl	Addition	No. pap 3 days	(papillae appear arrested in growth - failed to develop further after 2 days)
1	—	7		
2	0.1 M12 λ 25'	> 85		

16 papillae on 1 plate picked and streaked xyl - (see xyl-) 10/27 check to see if gal + - 10/29 see gal - ?

- 1924 t 1-12 #35 - this streaking = #4 from galdirection is mixed - no reisolation
Attempt to establish a stable + from unstable +.
12 well isolated + col. picked and streaked on EMB gal. Replicate 1

1	mixed
2	"
3	"
4	"
5	"
6	"
7	"
8	"
9	"
10	"
11 single colonies to base →	"
12	pure + → picked

- 1924 t 1-12 - One plaque type / 1485 - see 129
picked and streaked out EMB gal -

- 811 gal + reversions checked for their papillation behavior - on EMB base -

		Indication	
1307	5-10 pap/colony	=	
811	no. papillae		
811 gal + -1	< 1 pap/colony	mimic	does not grow on gal
2	< 1 pap/colony	"	" " "
3	< 1 pap/colony	"	
4	c. 5 pap/colony	true reversion	
5	5-10 pap/colony	" "	gal+ ! 811, 918
6	< 1 pap/colony	mimic	
7	c. 1 pap/colony	"	
8	5-10 pap/colony	true reversion	gal+ ! 811, 918
9	c. 1 pap/colony	mimic	
10	< 1 pap/colony	mimic	

Thursday 10/23/52

1924EK-12 #25 - from initial streaking for purpose of isolating stable + 4 streakings from pgs.
12 additional colonies streaked on EMB gal -

all mixed - many mutants - queer to EMC -

~

- Friday 10/24/52

1924EK-12 - the one plague type #5 of testing
streaked out EMB gal 10/22 -

~

1924EK-12 - #12 of Pg (131) for stability isolation
#25 colony
papillae

1st streaking from end of purification - pure
week 10/23 2nd streaking .. .. c. 150 colonies - all pure
10/24 3rd streaking .. .. c. 500 colonies - all pure
10/25 4th streaking .. .. c. 500 colonies - 1 (-) colony - contam.?

SEE PG 133

picked and Examined - see subsequent pg.

~

Saturday 10/25/52

1924EK-12 (above) 1 plague Type - transferred to agar slant (+ #1)
10 + and 10 (-) colonies picked from streaking and tested against 1405

+ colony	reaction	- colony	reaction
1	non lyr	1	swiger plague? → re-examined pg 134
2	"	2	"
3	"	3	"
4	"	4	"
5	"	5	"
6	"	6	"
7	"	7	"
8	"	8	"
9	"	9	"
10	"	10	"

Sunday 10/26

- 811 galt #5 lysate (from) - (this is papillating strain (on lawn)) - (0.5 ml of this lysate in Pa = sterile)

10^8 dil. = 44, 59 = 5.2×10^9 λ/ml

in 518 gives about 600 rp/0.1 ml = 6×10^3/ml $\frac{6 \times 10^3}{5.2 \times 10^9} = 1/10^6$

- 518 - Galdeletion

DNAase no effect

EMB gal	Addition	48 hrs	2 days
	none		(30)
2	0.1 K42λ (25') untreated *	960	
3	0.1 K42λ (25') treated **	998	
4	0.1 811 galt #5 λ	883	

some o plaques here?

gal reversion ↓ activity on gal⁻ →

examine for stability - see p 134

* untreated = 0.1 ml burst added to 0.5 ml lysate - incubated at room temp. 5 min

** treated = 0.1 ml DNAase added to 0.5 ml " - " " " "

"(1.1γ DNAase)

See also p. 135

DNA ase treated greatly decreased in viscosity

Boiled

~~~ 1929 EK42 #25 papillae col #12 - stable galt

over night culture (from) diluted plated EMB gal -

EMB Plate	no colonies	no. gal -
1	565	0
2	440	0
3	774	0
4	622	0
5	609	0
6	559	0
7	598	0
8	523	0
9	560	0
10	573	0
	5824	0

To here — Lusting

3 streaks mixed gal+, gal-
4 streaks gal+
1 plating of 5821 colonies no gal-
~~~

$\frac{2.1\times10^{3}}{5.2\times10^{9}} \cdot 0.5\times10^{-6}$

Tuesday 10/28

1821 gal- xyl-dilutions

EMB gal	Addition	No. papillae after 2 days
1	none	0
2	0.1 K-12λ (25')	4
3	0.1K-12λ (25')	7
4.	0.1K-12λ (25')	8

cultures for typ

- EMB xyl none
  - 1 ↙ 4
  - 2 ↙ 0
  - 3 ↙ 7 } faint background of many minute papillae c.10⁴

- 518t 811 gal + #5 ↗ pg 135

8 papillae picked and streaked - 4 papillae in the transduced of the type (1), colonies, red purple pigmented - papillae of same size (This is not unusual, reason for diff. not known) 4 red purple } streaked on EMB gal.
4 colonies (2)

probably not correct

1st	streaking -	all mixed +, -	no distinction as to above
2nd	" -	1/8 mixed +, -	mentioned above
3rd	"	1/8 mixed +, -	<- others giving off something intermediate

glucose fixed - this transduced site appears to be stable more quickly than 518 cells transduced by K-12 λ — this also appears to be the case for this lysate and 811 gal+ as below

- 1924 EC-12 One plaque type
  - recommunication of synicity of one plaque gal- type (p.135)
  - 4 gal- colonies from streaking examined - all non-synaict

- 8-12λ (X) 0.5 ml + 10 ml Ped Tested for sterility - sterile

- Wednesday 10/29/52
- 811 - gal- dilutions

EMB	Addition	No. pap. after 2 days
1	none	39
2	0.1 K-12λ (25')	196
3	0.1 811 gal+ #5 1 <- 1st lysate (viscous)	204

$\frac{1.57\times10^{3}}{1.6\times10^{10}} = \frac{1}{1.0\times10^{7}}$

<- stability of 12 of these examined

1st	streaking	7/11 mixed +, -
2nd	streaking	1/11 mixed +, -
3rd	streaking	1/11 mixed +, -

apparently → the activation is as with 518 t 811 gal + #5 above

This is probably. The case with 811 gal+ #8 on (p. 135a)

Thursday 10/30

1924tK-12S
Cross g- X Y-10

1. purity of 1924tK-12 culture -- well plated on EMB gel -

EMB gal	no. g+ col	no. g(-) col.
1	275	0
2	249	0
3	268	0
	792	

this are also additive ∴ data of p. 135 - make 5821 / 792 / 6613

2. Cross.

EMS gal	no. (+)	no. (-)
1	175 (1 mosaic)	0
2	116	1
3	152	0
4	141	0
5	165	—
	749	1

The 1 (-) and (1) mosaic colonies streaked out on EMB gal - both gave gal + [illegible] gal colonies

λ titrations.
grown on 1485
dil 10^6 = 0, 0, titer < 10^6

K-12 λ(X) in 240 seconds
dil 10^6 = 172, 262 = 217 = 2.2×10^9 ← previous assay says titer here originally was 2.8×10^9. Better recheck to observe if this post incubation assay is correct.

811 gal + #5 λ
untreated dil 10^8 = 37, 85 . 6.1×10^9 (previous p. 133 5.2×10^9)
DNAse treated dil 10^8 = 52, 68 = 6.0×10^9
} indicates DNA ase has little effect if any on either gal reduction or plaque production

DNA ase effect on plaque formation

efficacy of 811 gal + #5 λ = (for 518) $\frac{8.8 \times 10^3 \text{ gal/ml}}{6.1 \times 10^9 \lambda/\text{ml}} = 1.3 \times 10^{-6} = 1/7 \times 10^{-5}$

(for 811) $\frac{2.0 \times 10^3}{6.1 \times 10^9} = 0.3 \times 10^{-6} = 1/3.3 \times 10^{-6}$

811 gal reductions

	EMB gal	Addition		Papillae after 2 days
	1.	none		25
	2.	0.1 ml 811 gal + #5 1	untreated ≈ DNAse	201
	3.	0.1 ml 811 gal + #5 1	DNA ase treated	296
sterility of theta (0.5) ascertained	4.	0.1 ml 811 gal + #5 2	—	289 + ← plate contaminated?
	5.	0.1 ml 811 gal + #8 1	—	291 ←
	6.	0.1 ml K-12 (X) in 240		142

811 gal + #8 gal reduces!

518 gal reductions (control omitted)

EMB	Addition	Papillae after 2 days
1	0.1 ml K-12 (X) in 240 λ	540
2	0.1 (1/10) "	15 (dry plate) ← 10 examined / and 1 ↓ all gave faint mat / 1485
3	0.1 (1/100) "	9

(portrait)

Thursday 10/30

- 1402 galdeletion -

EMB gal	Addition	no plaq after 2 days
1	none	5
2	0.1 ml K-12 (25'), 1	43

} appears as a dilution in the absolute sense of the effect. Similar to result ~ 182/0.134 to this culture partially typ. R

811 gal + #1 X Y-10

- Cross for the purpose of determ. 811 gal + #1 suppressor or no deletion

EMB gal		no. gal (+)	no gal (-)
	1	333	0
(cumulative count) →	2	462	4
	3	541	2
	4	671	2
	5	781	1
		781	9 <- retest on EMB gal —

Summary table: (+) (-)
X1 gal 130 — 96
X2 gal 135a — 11 0
1126 1

- 811 gal + #2 X Y-10

- Cross for purpose of determ if this 811 deletion is suppressor mut. or no.

EMS gal		no gal (+)	no gal (-)
	1	157	0
cumulative count	2	326	2
	3	456	1
	4	603	0
	5	866	2
		866	5

} no gal slow -

Summary table: (+) (-)
X1 gal 122 — 0
X2 gal 123 — 0
X3 gal 126 275 0
X4 gal 135a 871 0
1879 0

- Sat. 11/1

811 + 811 gal + #8 stability ✓
1st streaking - 8/8 mixed +, -
2nd streaking - 0/8 mixed +, - (all +)
3rd streaking - 0/8 mixed +, - (all +)

- 2102 lysate for EMC -
mal. 35 sec - incubated 2 hours - centrifuged
dil 10^7 → against 2096 = no plaques - galdeletion on 518 according to EMC?

- 1673 mal for gal - on EMB gal
3 good gal mixes obtained - set for galdeletion

1673 gal - #1 #2 #3

- 1692 mal. for gal - unsuccessful attempt

Sunday 11/2

- 1655 X 1177 to obtain xyl- for xyl distrib attempt
  old suspension used - small number of prototrophs
  c. 6 mult from c 100 prototrophs.
  picked and streaked on EMB xyl. — 4/6 xyl minus
  pick and see if λ^s - streak on other med. —

- 1924tK-12 #12 (unstable) X 1436 - to see if λ^+ obtained — but the dmn. to λ^+ cross should give λ^+ if this is so since λ^r recessive to λ^+. Examine unstable gal+ prototrophs to see λ^+ vs since $\frac{gal\ \lambda}{+\ +}$ should go together

1924tK-12 parent culture - Analysis - streaks of washed susp. made / 1485 - no evidence of λ

Gal+	Plate	#(+)	#(-)
	1	$\frac{1}{4}$ = 200 (800)	$\frac{1}{4}$ = 11 (44)
	2	$\frac{1}{4}$ = 161 (644)	$\frac{1}{4}$ = 9 (36)
	3	$\frac{1}{4}$ = 173 (692)	$\frac{1}{4}$ = 8 (34)
		2136	112

} purity of 1924tK-12 parent

Cross	Plate	#(+)	#(-)
	1	42	383
	2	38	354
	3	—	—
	4	..	..
	5	68	365
		148	1102

about the same as first two but difficult to count due to agar seed background

28+ picked and streaked against 1485- all non lysogenic

Thursday 11/6

- 1673 gal- fractures

		EMB gal	Additions	No gap. after adapt
#1 (no 1 and 2 gal. plates inferred)	1	0	none	7
	2		0.1 ml 1K-12 λ (25')	$\frac{1}{8}$ = 195 = 1560
#2	1		none	11
	2		0.1 ml 1K-12 λ (25')	$\frac{1}{8}$ = 163 = 1304
#3	1		none	19
	2		0.1 ml 1K-12 λ (25')	$\frac{1}{8}$ = 161 = 1288

10/6 Thursday

– 1436. galduction to Hfr unstable + which is λ^+ gal$^+$ S^R
to cross with 578 to observe if lysogeneity of galduced is due to donor nonce of λ^+ in transduced fragment → examine to see if unstable i.e. S^a - possibility of transducing S^s which is dominant



1. galduction of 1436

	EMB gal	Additions	Papillae after 2 days
	1	none	34
	2	.01 ml K+12λ (25')	1/8 = 136 = 1248

what kind of prototrophs?
gal+ $\lambda^?$ $S^?$
gal– $\lambda^?$ $S^?$
design control
test examine whose problem

① ← 1st Pure. streak 7/8 gal– ! what ? SEE 112
② try again 1st Pure. streak 7/8 gal+
③ try again 1st pure. streak 15/18 gal– of this lot 9 streaked against 1445 all lysogenic

Pap #	Streak 1	Streak 2
1	mixed	pure +
2	mixed	"
3	mixed	"
4	gal –	—
5	mixed	pure +
6	mixed	"
7	mixed	"
8	mixed	"

207° = 673 gal– –1. Stability check –

Sputum	1st Streak	2nd Streak	3rd Streak
1	mix	pure +	pure +
2	pure +	"	"
3	pure +	"	"
4	mixed	"	"
5	pure +	"	"
6	pure +	"	"
7	pure +	"	"

207° ✓

Galduced	1st Streak	2nd Streak	3rd Streak
1	mix	mix ⊗	mix ⊗
2	mix ⊗	mix ⊗⊗⊗	" ⊗
3	mix	mix ⊗	" ⊗
4	mix ⊗⊗⊗	mix ⊗	"
5	mix ⊗⊗⊗	mix ⊗	"
6	mix	pure +	pure +(?)
7	mix ⊗⊗⊗	pure +	" +

673 gal– –2

1	mix	pure +	pure +
2	mix	mix	"
3	pure +	pure +	"
4	mix	"	"
5	mix	"	"
6	pure +	"	"
7	mix	"	"
8	mix	"	"

1	mix	mix ⊗	pure +
2	pure +	mix ⊗	pure +
3	mix	mix(?)	pure +
4	mix	mix	mix ⊗
5	pure +	pure +	pure +
6	pure +	mix ⊗	pure +
7	mix	pure +	pure +
8	mix	pure ⊗	mix ⊗

673 gal– –3

1	mix	pure +	pure +
2	mix	"	pure –
3	mix	"	pure +
4	mix	"	"
5	mix	"	"
6	mix	"	"
7	mix	"	"
8	mix	"	"

1	mix	pure +	mix ⊗
2	mix ⊗	pure +	mix ⊗
3	mix	pure +	pure +
4	mix ⊗	mix	mix ⊗
5	pure +	pure +	pure +(!)
6	mix	pure +	mix ⊗
7	mix	pure +	pure +
8	mix ⊗	mixed ⊗	" +

Sunday-Monday 18/9/18/10

- ~~W1436 [illegible]~~

- 1177X 1655 xyl- #1

EMB gel	Addition	No pap 2 days
1	none	0
2	0.1 ml K12 λ (1')	0

Attempted xyl induction of 1177 xyl-

#2		
1	none	0
2	0.1 ml K12 λ (2')	0
#3		
1	..	0
2	..	0

811 transduction ~ 750 λ

EMB gel	Addition	No pap. 2days
1	none	45
2	0.1 ml K12 λ (2')	55

small increase - xyl? pick a few - look for motability

slightly large [illegible] pap. Spread top small [illegible]

Select to do?

818 transduction ~ 750

EMB gel	Addition	No pap 2d
1	none 750 λ	45
2	0.1 ml K12 λ (1')	40

818 p2R = W1412

EMB gel	Addition	# pap. 2d
1	none	33
2	0.1 ml K12 λ (15')	35

818 p.R not gal-mob

Sunday 11/16

1673 gal- #2 × Y-10 – as usual – no control –

EMB gal	Lac+	Lac−
1	324	12
2	166	7
3	261	10
	751	29

(+) slightly purple – some (−) mosaic

← 26 picked EMB gal, 22 (−) streaked /1485/λ SEE BELOW. – 7/22 Sequestrum

≈ 30 picked EMB gal – streaked against /1485, λ 30/30 lysogenic on 1485

1673 gal #3 × Y-10 as above

EMS gal		
1	174	16
2	186	13
3	418	20

(−) slightly purple – some (+) mosaic

518 gal mixed ē 892 λ Gal3- mixture on Gal4- 1S

6 picked streaked EMB gal

	strk 1	strk 2	strk 3
1.	mix	mix	mix – STOCK 518t892-1
2.	pure+(?)	pure+	—
3.	pure+	pure+	—
4.	pure+	pure+	—
5.	pure+	pure+	—
6.	mix	mix	mix STOCK 518t892-2

518t892

EMB gal	Addition	No pfp. 2 days
1	none	41
2	0.1 892 λ	247

Product

811 gal mixed ē 892 λ mixture Gal3- on Gal4- λt

6 picked pure streaked EMB gal

	strk 1	strk 2	strk 3
1	mix	pure+	
2	"	mix	mix
3	"	mix	mix
4	"	mix	mix
5	"	pure+	—
6	"	pure+	—

811t892

1	none	39
2	0.1 892 λ	85

$\frac{4.6\times10^{2}}{3.9\times10^{9}} = \frac{1}{8.5\times10^{6}}$

STOCKS 811t892-1, 811t892-2, 811t892-3

Tuesday 11/18

1673 gal-#2 XY-10 above gal- prototrophs tested –

			1485	λ			1485	λ
Lp+	1	○	lys-	resist	Lp+	12	○	lys	resist
Lp+	2	○	"	"	Lp+	13	○	"	"
Lp^s	3	○	non lys	sensitive	Lp+	14	○	"	"
Lp+	4	●	lysogenic	sensitive	Lp^s	15	●	Sglegma	~~sensitive~~ sens.
Lp+	5	●	lysogenic	sensitive	Lp+	16	○	lys-	resist
Lp^s	6	○	non lys	~~sensitive~~ sens.	Lp+	17	○	"	"
Lp+	7	○	lys	resist	Lp+	18	○	lys	resist
Lp^s	8	◐	non lys	sens	Lp+	19	○	lys	resist
Lp+	9	○	lys	resist	Lp^s	20	○	~~non~~ lys	sensitive
Lp+	10	○	lys	resist	Lp^s	21	○	~~non~~ lys	~~sensitive~~ resist
Lp	11	◐	single plaque	sensitive	Lp+	22	○	lys	resist

gal- Lp+ 15
gal- Lp^s 7

replica to D(5)

Lp+ and Lp^s prototrophs

all prototrophs

Wednesday - 11/19

- S1Bt transduction by 902λ - obtain stocks

EMBgal	additions	no. pap. 2 days
1	none	39
2	0.1 902λ	1/4 = 288 1152

- 811 transduction by 902λ

EMBgal	additions	no. pap 2 days
1	none	44
2	0.1 902λ	147

$\frac{1.03 \times 10^3}{4.9 \times 10^{10}} = \frac{1}{4.9 \times 10^6}$

Strains made S1Bt 902-1, -2

	1.	2.	3.	4.	5.	6.	7.	8.
	mixed	"	"	"	"	mixed	pure +	mixed
	mixed	"	"	"	"	pure +	pure +	mixed
	mixed	"	"	"	"	pure +	mixed	mixed

Exp.	1st streak	2nd streak	3rd streak
1.	mixed	mixed	mixed
2.	"	"	"
3.	"	"	"
4.	"	"	"
5.	"	"	"
6.	"	"	"
7.	"	"	"
8.	"	"	"

- 1436 (purified) Recheck on gal- streaking - 9 deduction

EMBgal	additions	no. pap 2 days
1	none	6
2	0.1 IC-12 λ (25')	1/4 = 294 1176

← sixteen (16) picked and streaked - all gal-, all non-protomorphic

Thursday 11/20

- S1BtK-12 X Y-10 - for the purpose of uncovering gal hp^s?

	EMBgal	no +	no -	no. mixed
S1BtK-12 purity ✓	1	384	72	8
	2	406	60	9

EMSgal		
1	184	2 —
2	172	3 —

4 examined all λ^R all lysogenic on 1485

Booked

- 811K-12 X 1673 - for the purpose of uncovering (gal- hp+)
X 1678

	EMBgal	no(+)	no(-)	no mixed
811BtK-12 purity ✓	1	551	43	8
	2	633	37	4

X1673 EMSgal		5	—
1	73	2	—
2	91		—

7 examined - all λ^R all lysogenic on 1485

X1678 EMSgal		2	
1	101 contam.	4	
✓	56		

5 examined all λ^R all lysogenic on 1485

Thursday 11/20

Friday 11/21

- 1673-gal- -3 X Y- 10 - [illegible] 139 - lysate aggregating

Check on λ-state of protohaploids

gal-	1.	lys	R	14	
	2	lys	R	15	
	3	lys	R	16	
	4	lys	R	17	
	5	lys	R	18	
	6	lys	R	19	
	7	lys	R	20	
	8	lys	R	21	
	9	lys	R	22	
	10	lys	R	23	
	11	lysect	R	24	
	12	[illegible]	S	25	
	13	lys	R	26	



1485	λ
no plaque	R?
lys	R
no lys	sens
lys	R
lys	sens ?
lys	R
no	sens ?
lys	R
no lys	sens
lys	R
lys	R
lys	R

10 (-)

gal+ 22 picked all lys on 1405, all λ resist

Salmonella for JL.

- lysate attempted in 914, 915, 922

lysates in 914-1 some suggestion of lysis?
913 — discarded

lysate in 914-2
915
922

Results

914-1 / 913 → cross streaks on 913 - ① - activity in undiluted
914-2 / ② " " undiluted
known suppressed (JL) " " "

915 / 912 → cross streaks on 912 ① (915) - activity undil., some act. in 10^6 dil? hopeful

② (922) - activity in undil, none in 10^6 -

(142) 152/16 = 9 17/2 / 2432

Tuesday 11/26

– New λ preparation.

518 (examination of allogenic ~~[illegible]~~ gal+ as donors of gal+) –

EMB gal	Addition	No papillae 2 days
1.	none	28
2.	0.1 518tC8g2λ-1	3
3.	0.1 811tC8g2λ-1	0
4.	0.1 K-12λ (11/23)	1/16 = 152 = 2432

} recently observed by numerous investigators

borrowed

– Other gal⁻ mutations – examination for transduction –

stock mode of plaques

	EMB gal	Addition	No papillae 2 days
566 ← (tp2 1)	1	none	9
	2	0.1 K-12λ (Y)	9
583	1	none	6
	2	0.1 K-12λ (Y)	340

picked	streaking 1st	streaking 2nd	streaking 3rd
1.	mixed	mixed	-
2.	"	"	"
3.	"	"	"
4.	"	"	"
5.	"	"	"
6.	"	"	"
7.	"	"	"
8.	"	"	"

– 2050 gal3 – Examination of allogenic transduction (lysates) for activity on gal3

EMB gal	Addition	No papillae 2 days	
1.	none	29	
2.	0.1 518tC8g2-1λ	solid smear	(3 papillae?) ← thin lysate not sterile
3.	0.1 811tC8g2-1λ	solid smear	(3 papillae?) thin lysate not sterile
4.	0.1 811λ	450	

2050tC811 stock →

$\frac{1.7\times10^{10}}{4.3\times10^{3}} = \frac{1}{4.0\times10^{6}}$

see 144 – 518tK-12 – isolation of several gal⁻ segregants to see if

1. gain no gal+ ε X 1436
2. all lysogenic
3. X 1673 to see if all gal⁻ tp⁺ (linkage intact)

a. streak out overnight culture 518tK-12

– 518tK-12 – crosses with 1678 gal+tp⁺tp^s

gal+ tp^s / gal⁻ tp⁺ –

	EMB gal	Condition	(+)	(–)	
borrowed	1	0.1 ml of each overnight cultures	6	8	B
	2	" " " "	11	6 (+)	B
518tK-F⁻	3	Aerated 518tK-12, unaerated 1678	0	0	what happen?
	4	" " "	0	0	
	5	" " "	1	0	
	6	" " "	0	0	
	7	" " "	heavily contam	0 (–)	
1678 F⁻	8	Aerated 1678, unaerated 518tK-12	78	2/16	Some gal+ appear to be mixed
	9	" " " "	70	30	
	10	" " " "	72	18	
	11	" " " "	76	26	
	12	" " " "	83	25	

of the 115 – 110 streaked single colony picked and tested against 3 – all 1R – replica'd against 1485 – all lysogenic

(+) (–) individual strokes – A

Saturday 11/29

— Reexamination of the conjugability of aerated 518tK12 cultures —

1. 518(~~streptomycin~~) X 1678, 1673 — (all cultures resuspended in 1X Saline)

EMB gal	Additions	(+)	Petri dish (-)	
1	aerated 518, unaerated 1678	0	18	F+1678 XF-518 = 0/18
2	" " 1678	0/0	53	F-1678 XF+518 = 0/53
3	aerated 518, unaerated 1673	0	10	F+1678 XF-518 = 0/71
4	" " " "	1/1	18/28	F-1678 XF+518 = 3/57
5	unaerated 518, aerated 1678	3	571	F+1673 XF-518 = 17/28
6	" " " 1673	16	411	F-1673 XF+518 = 16/427

(booked)

2. 518tK12 X 1678, 1673

7	aerated 518tK12, unaerated 1678	0	1	
8	" " " "	1	0	
9	aerated 518tK12, unaerated 1673	0	0	
10	" " " "	0	0	
11	unaerated 518tK12, aerated 1678	135	28	← (of these 28 I examined - all λ^R all lys / 1485)
12	" " " 1673	146	32	

(booked)

— Sterility checks

518t892-2	1.0 ml
518t892-1	1.0 ml
811t892-1	1.0 ml
811t892-2	1.0 ml

in Penassay - o.k.

-Sunday 11/30

2050 Reexamination of putative C 2050 in ~~Fig.~~ 142

EMB gal	Additions	24 hrs 2 days
1	none	½ = 131 (what happened?) (all small)
2	0.1 518t892-1	solid growth gap (mostly +?) c. 5000?
3	0.1 518t892-2 (undiluted)	solid growth gap (mostly +?) c. 5000?
4.	0.1 518t892-2 (dil 1-10)	solid growth gap (mostly +) ½ = c. 100 = 7000
5.	0.1 811t892-1 (und)	solid growth gap (mostly +) c.
6.	0.1 811t892-1 (1-10)	solid growth gap (mostly +)
7.	0.1 811t892-2 (und)	½ = 94 (all small
8.	0.1 K-12 (11/13) λ	½ = c. 100 = 800



No papillae →

A1 wo = wo and 573 gal
A1 K-12 = 0.1 K-12 λ " "
A1 891 = 0.1 892 λ " "

Monday-Tuesday 12/1 12/2

5% - No of cells plated - influence on number of transductions observed (papulon)

Stock suspension dil = 2×10^7 — 648,547 = $6.97\times2\times10^7$ = 139.4×10^7 = 1.4×10^{10} cells/ml
dil. stock plated in constant, 0.1 ml K-12 λ (11/23)

# cells	used	in K-12 λ lysate (2504)	in at K-12 λ
1.4×10^8			
7×10^8	1-2	½=445 (1780)	35
3.5×10^8	1-4	¼=369 (1472)	29
1.75×10^8	1-8	⅛=280 (1120)	10 *
8.75×10^7	1-16	1/16=662 (1324)	33
4.4×10^7	1-32	1/32=581 (1862)	49
2.2×10^7	1-64	1/64=251 (562)	39
1.1×10^7	1-128	535	32
5.5×10^6	1-256	509	23

(plotted)

titer 2.7×10^{10} (estimated)

* heavily contaminated ∴ spore plaques

14% transduced by 750 - Distinction between gal$_1$- and gal$_4$-

EMB gal	Addition	No pap. 2 days
1	none	24
2	01 750 λ	145

← many of these papillae have haloes about them spontaneous do not.

16 papillae picked from EMB gal 1 streaked on gal

	1st alt (-)	2nd alt (-)	3rd alt (+)		1st alt (-)	2nd alt (-)	3rd alt (+)
1	..	..	..	9	..	..	..
2	..	..	..	10	..	..	..
3	..	..	..	11	..	..	..
4	..	..	..	12	..	..	..
5	..	..	..	13	..	..	..
6	..	..	..	14	..	..	..
7	..	..	..	15	..	..	..
8	..	..	..	16	..	..	..

discarded

2050 ± 811 pfu (142) stability

8 picked	1st mixed	2nd mixed pure (+)*	3rd mixed? pure (+) (+)
1.	..	..	..
2.	..	..	..
3.	..	..	..
4.	..	..	..
5.	..	..	..
6.	..	..	..
7.	..	..	..
8.	..	..	..

← Stock made of this

* + reaction not as strong as usual-

Sunday 12/7/52

5784K-12 · Examination of gal- segregants for their allelic state · (–) derived from pectored gal+ colonies · 6 streaked out once and single (–) colony inoculated into Pen. Overnight cultures [illegible] as follows:

Strain/gal(–) /1485		No add (contaminated ē (+) colonies) →	K-12 (λ)	811-2 λ	902 λ-1	Conclusion
	1					
	2	23	130 (224)	14	262	gal+
Gal– segregants from 5784K-12	3	(contaminated ē (+) colonies) →				—
	4	(contaminated ē (+) colonies) →				
	5	23	98 (225)	24	231	gal+
	6	(contaminated ē (+) colonies) →				—
	7	29	226 (273)	28	302	gal+
	8	(contaminated ē (+) colonies) →				
	9	31	146 (216)	30	248	gal+
?	10	48	186 (225)	80	299	gal+ (?)
	11	37	154	22	284	gal+
	12	40	164	53	245	gal+
This series = A	13	(contaminated ē (+) colonies) →				—
	14	32	103	22	263	gal+
	15	41	84	23	345	gal+
	16	28	138	20	271	gal+
	17	30	87	27	287	gal+
	18	20	83+	24	231	gal+
	19	28	67	28	239	gal+
	20	28	107	34	206	gal+
	21	31	97	17	245	gal+

↖ magnitude of 902-deletion similar to that of 902 on 811 which with this prep was (147±202) - c 40

16/17 say (–) segregants are gal+

1/17 says (–) segregant is gal+ - with possible change in ct.

now 17/17 say gal+

this 811 on lysate

EMB	Added	Pop 2 days
1.	811 cell - no λ	48
2.	" (811 λ)-2	50
3.	902 x 1658 lysate - no λ	13
4.	" - (811 λ)-2	105
5.	2050 - no λ	42
6.	" - (811 λ)-2	289

Exam. 811 plate stable

Strain	1	2	3
1.	mixed	mixed	
2.	"	"	
3.	"	"	
4.	"	"	
5.	"	"	
6.	"	"	
7.	"	"	
8.	"	"	
9.	"	"	
10.	"	"	
11.	"	"	
12.	mixed	mixed	
13.	"	"	
14.	"	"	
15.	"	"	
16.	"	"	

5B Note. h-10 unstable, revertants EMB gal+ none = 74 2 (0.5B λ)-2 = 59 apparently 1st assay in error - contam ē spreader

4 failures

17 successful tests

2070 = (1673 gal– 1) examined for its relationship to gal 1-4 – by means of lysate activity.

EMB gal	Addition	No. prop. 2 days
1	none	10
2	0.1 750 λ	54
3	0.1 902 λ	¼ = 314 1256
4	0.1 892 λ	331
5	0.1 811 λ	175

2070 t 750

	1	2nd mixed	2nd mixed
1	mixed	mixed	mixed
2	+	+	+
3	+	+	+
4	+	+	+
5	+	+	+
6	+	+	+
7	+	+	+
8	mixed	mixed	mixed

2/4

2070 t 892

	1	2	3
1	mixed	+	+
2	+	+	+
3	+	+	+
4	+	+	+
5	+	+	+
6	mixed	mixed	mixed
7	mixed	mixed	mixed
8	mixed	mixed	[illegible]

2/4

2070 t 902

	1	2	3
1	mixed	mixed	mixed
2	+	+	+
3	mixed	mixed	mixed
4	mixed	mixed	mixed
5	mixed	mixed	mixed
6	+	+	+
7	+	+	+
8	mixed	mixed	mixed

5/8

2070 t 811

	1	2	3
1	+	+	+
2	+	+	+
3	+	+	+
4	mixed	+	+
5	+	+	+
6	mixed	+	+
7	+	+	+
8	+	+	+

0/8

transferred to ē plate for

518+902 gal-
neg
= B transduce ε 811
902

12/9/52
Tuesday

146

518+902 Gal- segregants - Examination - Isolated from single segregating colonies streaked on EMB plates - tested a Bacteriophage disease

	1485	no. spots			gal+
1.	lysogenic	235	18	50	gal+
2.	"	223+	29	16	4-
3.	"	102+	22	15	4-
4.	"	contam.	29	24	4- ?
5.	-	137+	18	14	4-
6.	"	143+	25	23	4-
7.	"	10	507	9	2- ← }
8.	-	10	474	7	2- }
9.	"	325+	191	214	4-
10.	-	151+	16	31	4-
11.	-	160	20	44	4-
12.	"	232+	44	52	4-
13.	"	143+	16	32	4-
14.	"	136+	18	36	4-
15.	-	46	32	~~38~~ 38	? }
16.	-	116+	29	21	4-
17.	-	214	31	31	4-
18.	"	99+	30	33	4-
19.	"	139+	41	41	4-
21.	-	contaminated ε gal+			
21.	"	22+	21	23	4-
	streaked	0.1 ml	0.1 ml	no add	
	1485	902 λ	(811 λ)-2		
					17 4-

17 gal4-
2 gal2-
1 double?

The ocrease - B

Isolated

Saturday 12/13

C. -578 & 892 Examination of gal- segregants

gal- isolate	1485 lysogenic	0.1 (892 II)-1	0.2 (811 A)-2	[illegible]	probable locus
1.		92	22	24	gal 4 -
2.	"	78	38	36	gal 4 -
3.	"	99	27	25	gal 4 -
4.	"	353	304	20	*
5.	"	84	26	18	gal 4 -
6.	"	68	29	19	gal 4 -
7.	"	contaminated c + cells →			
8.	"	376	392	29	*
9.	"	442	324	45	*
10.	"	453	410	21	*
11.	"	337	364	30	*
12.	"	99	25	31	gal 4 -
13.	"	427	294	46	*
14.	"	387	416	23	*
15	"	78	36	32	gal 4 -
16	"	contaminated c + cells →			
17.	"	78	30	11 (mid plate)	gal 4 -
18.	"	309	271	22	*
19.	"	296	257	3	*
20.	"	88	25	35	gal 4 -
21.	"	73	29	38	gal 4 -
22.	"	67	33	25	gal 4 -
23.	"	278	190	21	*
24.	"	309	207 (thin plate)	18	*

booking

NG

892 a

mut

11 gal 4 -
* 11 Unknown locus -

strains

Stability examination of * mutants locus -

	C-4-811			C-4-892			C-4-811			C-4-892		
	①	②	③	①	②	③	①	②	③	①	②	③
1.	mixed	mixed	mixed	mixed	mixed	mixed	mixed	+	mixed	mixed	mixed	mixed
2.	mixed	mixed	-	-	-	-	-	mixed	-	-	-	-
3.	mixed	mixed	-	-	-	-	-	-	-	-	-	-
4.	+	+	-	-	-	-	-	-	-	-	-	+
5.	+	-	-	-	-	-	-	-	-	-	-	mixed
6	mixed	-	-	-	-	-	-	-	-	-	-	-
7.	mixed	-	-	-	-	-	-	+	-	-	-	-
8	mixed	-	-	-	-	-	-	mixed	-	-	-	-

Worked

All galR cultures transferred to a plate for stock purposes

Saturday 12/13

S18TK42 - V on stability of gal-segregant after a second transduction - See pg 145 repeat exp

gal-pplate	1st streak	2nd streak	3rd streak
A-2 pp 1.	+	+	mixed
2.	mixed?	mixed	mixed
3.	+	+	+
4.	+ slow growth?	mixed	+
5.	+	+	mixed
6.	+	+	mixed
7.	+ slow growth?	+	+
8.	+ 1/8	+ 2/8	+ 2/8
A-5 pp 1.	mixed	mixed	+
2.	+	mixed	mixed
3.	+	+	+
4.	mixed	+	mixed
5.	mixed	mixed	+
6.	+	mixed	+
7.	+	mixed?	+
8.	mixed 4/8	+ 6/8	+ 7/8
A-7 pp 1.	mixed	mixed	mixed
2.	"	mixed	mixed
3.	"	mixed	mixed
4.	"	+	+
5.	"	+	+
6.	+	mixed	mixed
7.	+	+	mixed
8.	+ 1/8	mixed 5/8	mixed 6/8
A-9 pp 1.	mixed	mixed	mixed
2.	+	mixed	mixed
3.	mixed	+	+
4.	+	mixed	mixed
5.	+	+	mixed
6.	mixed	+	+
7.	+	mixed	mixed
8.	mixed 4/8	mixed 5/8	mixed 6/8
A-11 pp 1.	+	+	mixed
2.	mixed	mixed	mixed
3.	mixed	mixed	mixed
4.	mixed	+	+
5.	+	+	+
6.	+	mixed	mixed
7.	mixed	mixed	+
8.	+ 4/8	mixed 5/8	mixed 5/8

Bottled

previously (p.96) → 4 transfers = 60% mixed
here 23/40 → 3 transfers = 58% mixed

0.57 = 58
40 | 23.0
200
300
280
200

Monday 12/14

Crosses.

811 XY-10 to obtain gal- prototroph to examine the ability of 811 gt-1 and 811-gt-2 to transduce in minimal - Do the weak gal+ of these reversants another allele not having sufficient power to allow for selection in EMB gal ?

		prototroph. (+)	(-)	
EMB gal 1	both Y-10 × 811	237	8	
2.	" "	226	9	
		463	17	17/480

811 gt-6 XY-10 to observe if this is true reversion

	(+)	(-)	
EMB gal 1. both.	277	0	
2. "	357	0	
3.	341	1	← in clones on EMB gal this colony +
	975	1	

811 gt-7 X Y-10

1.	both	145	0
2.	"	215	1?
3	"	254	0
		614	1? ∴ ← This col also + on ✓

811 gt-9 X Y-10

1.	both	235	0
2.	"	245	1?
3.	"	280	1?
		760	2? ∴ ← both + on ✓

811 gt-10 X Y-10

1.	both	205	0
2.	"	153	0
3	"	205	0
		563	0

2062
Natural λ

EMB gal	no. pos 3 days
1 no add'd	4
2 a) 1485 λ-1	12
3 0.1 K12 λ (11/23)	219
0(a) 1. no add	0
2. a) 1485 λ	0

Lysate sterility

0.1 of 902, 892, 750 λ into Pen - all sterile -

Monday 12/18/52

518 transduced gal- segregants.

B = 518 t 902

B7 = on prelim. test, gal$_2$ -

EMBgal		after 2 days
1.	no add	56
2.	0.1 (650 λ) -1	611
3	0.1 (902 λ) +1	58
4.	0.1 (892 λ) -1	342
5.	0.1 (811 λ) -3	147 ← this prep contam.

12/23

	Keap't dunks
	63
	463
	65
	387
← this prep contam.	205 ← (811 λ) -3

Crosses

	(+)	(-)
B7 X 902 -	4	390
"	7	277
	13	667 + 225 = 892

↑ not releasable 902 contam. E +

B-8 = gal$_2$ - on prelim exam.

EMBgal		
1.	no add	59
2.	0.1 750 λ -1	705
3.	0.1 902 λ -1	77
4.	0.1 892 λ -1	513
5.	0.1 (811 λ) -3	173 ← this prep contam.

despite this it apparently has some activity - other case contam. sometimes showing up in B-15 as well - duckweed

Crosses

B-8 X 902	1	137
	3	176
	1	145
	5	458

this cross repeated giving about 12 <1% +. Discarded by accident - involved about 10 plates with c. 200 prototrophs / plate

B-15 = a double ? gal$_2$ - gal$_4$ - ?

EMBgal			
1	no add	30	(many small -)
2.	0.1 (650 λ) -1	35	(" ")
3	0.1 (902 λ) -1	36	" "
4	0.1 892 λ -1	27	" "
5.	0.1 (811 λ) -3	28	" " ← this prep contam.

Crosses

B15 X 902	3	118
2.	3	176
	6	294
B15 X 892	1 ?	5
	8	4
	1 ?	9
B-15 X 1476	0	322

either gal$_2$?

C = 518 t 892

C4 = transduced by 811 and 892 - a new allele?

EMBgal		
1.	no add	46
2.	0.1 (750 λ) -1	286 (wet plate)
3.	0.1 (902 λ) -1	27
4.	0.1 (892 λ) -1	190 (wet plate)
5.	0.1 (811 λ) -3	83 ← this gal$_4$ contam.

Crosses

C-4 X 902	2 ?	174
	3	160
	5 ?	334
C4 X 892	1	10
	1	6
	2	16
C4 X 1476	0	155
	0	129
	0	284

probably caused by contam in 902 culture

C9 = as C4

EMBgal		
1.	no add	72
2.	0.1 (750 λ) -1	586
3	0.1 (902 λ) -1	91
4.	0.1 (892 λ) -1	397
5	0.1 (811 λ) -3	190 ← this gal$_4$ contam.

Crosses

C-9 X 902	1	165
	0	132
	1	297
C-9 X 892	0	0
	1	5
	0	5
C9 X 1476	0	108
	1	117

boosted

boosted