

CHAPTER 1

	Page
CARDIOVASCULAR DISEASES	
CORONARY HEART DISEASE (CHD)	
<i>Introduction</i>	3
<i>Epidemiologic and Autopsy Studies</i>	4
I. Studies in Men Demonstrating a Relationship Between CHD and Smoking	4
II. Studies Failing to Demonstrate a Definite Relationship Between Smoking and CHD	8
III. Studies in Women Relating CHD and Smoking	9
Carbon Monoxide (CO)	
<i>Introduction</i>	10
<i>Epidemiologic Studies</i>	11
<i>Experimental Studies in Man</i>	11
Nicotine	13
<i>Experimental Studies</i>	13
PERIPHERAL VASCULAR DISEASE	14
<i>Epidemiologic Studies</i>	14
<i>Experimental Studies</i>	16
CEREBROVASCULAR DISEASE (CVD)	16
<i>Epidemiologic Studies</i>	16
CIGARETTE SMOKING AND ASSOCIATED CHD RISK FACTORS	17
THROMBOSIS	18
SUMMARY OF RECENT CARDIOVASCULAR FINDINGS	19
CARDIOVASCULAR DISEASE REFERENCES	19
CARDIOVASCULAR DISEASE SUPPLEMENTAL REFERENCES	22

List of Figures

Figure 1.—Mortality for men with and without ventricular premature beats (VPB), according to presence or absence of smoking as a risk factor. Bars show percentages of men who died during 3-year period	5
Figure 2.—Relative odds of developing atherothrombotic brain infarction (ABI), coronary heart disease (CHD), or intermittent claudication (IC) according to levels of cigarette smoking	15
Figure 3.—Serum cholesterol levels of cigarette smokers and nonsmokers in the Belgian military service	18

List of Tables

Table 1.—1958 serum cholesterol, diastolic blood pressure, cigarette smoking status, and 12-year mortality—cohort of 903 men, age 40 to 59 in 1958, free of evidence of organ system disease and ECG abnormalities and followed long-term without systematic intervention. Peoples Gas Company Study, 1958 to 1970	7
Table 2.—Cigarette smoking and sudden death from CHD in women	10
Table 3.—Standardized multivariate regression coefficients of intermittent claudication on various characteristics	15

CHAPTER 1

CARDIOVASCULAR DISEASES

CORONARY HEART DISEASE (CHD)

Introduction

One million deaths per year are attributable to diseases of the cardiovascular system in the United States. Arteriosclerotic cardiovascular disease (ASCVD) is the leading cause of death in this country, accounting for greater than 50 percent of annual deaths; coronary heart disease (CHD) alone is responsible for 600,000 deaths per year. Cigarette smoking, hypercholesterolemia, and hypertension have been identified as major risk factors for the development of CHD.

Epidemiologic, autopsy, and experimental evidence presented in past editions of this report (1964, 1967, 1968, 1969, 1971, 1972, 1973) support a causal relationship between cigarette smoking and cardiovascular morbidity and mortality, as summarized below:

1. Both retrospective and prospective epidemiologic studies have demonstrated a strong relationship between cigarette smoking and increased CHD morbidity and mortality, with approximately a twofold higher risk of dying from CHD for all male cigarette smokers compared to nonsmokers.

2. A dose-response relationship has been demonstrated between cigarette smoking and CHD morbidity and mortality in men.

3. Cigarette smoking acts both independently of and synergistically with the other two major risk factors to produce these effects on CHD morbidity and mortality.

4. The above relationships between cigarette smoking and CHD morbidity and mortality have been demonstrated in Black and Asian, as well as Caucasian, populations.

5. The relative importance of cigarette smoking in the development of CHD in young men (less than 50 years old) is greater than that for any other risk factor.

6. Most prospective and retrospective studies suggest that pipe and cigar smokers exhibit a slightly higher risk of development of CHD than nonsmokers (but a significantly lower risk than cigarette smokers), while some studies demonstrate no such relationship.

7. Prospective epidemiologic studies document that cessation of cigarette smoking results in reduced mortality from CHD.

8. Autopsy studies reveal greater frequency and severity of coronary and aortic atherosclerosis among cigarette smokers than nonsmokers; cigarette smokers have been reported to have greater myocardial arteriolar wall thickening at autopsy than nonsmokers.

9. Experimental evidence in humans suggests that cigarette smokers with preexistent angina have a greater impairment in cardiac work capacity than nonsmokers. The role of cigarette smoking in the etiology of angina is unclear.

10. Experimental studies on humans and animals have shown that the pathophysiologic changes commonly observed in patients with CHD may be aggravated by cigarette smoking; contributions from both nicotine and carbon monoxide have been demonstrated. In addition, some of the biochemical and anatomical abnormalities seen in CHD have been induced by cigarette smoke, carbon monoxide, and nicotine.

Most of the studies reviewed in the last year confirmed the knowledge of the relationship between cigarette smoking and CHD. A listing of these studies appears in a separate section of the Supplemental Bibliography. A number of studies extended the knowledge of the association between cigarette smoking and CHD, but several studies presented data which were either partially or wholly inconsistent with the known relationships; these two types of studies are reviewed below.

Epidemiologic and Autopsy Studies

I. Studies in Men Demonstrating a Relationship Between CHD and Smoking.

The Coronary Drug Project Research Group studied, by multivariate analysis, 2,035 survivors of myocardial infarction in a prospective investigation to determine the relationship of placebo-treated premature beats and other known risk factors to sudden coronary deaths and total mortality (CV 12). Nonsmokers with ventricular premature beats (VPB) had a significantly higher total mortality than nonsmokers without VPBs ($P < .05$), and smokers with VPBs had a higher incidence of sudden coronary

deaths than smokers without VPBs ($P < .01$). Among patients without VPBs there was a significantly higher mortality rate in smokers compared to nonsmokers ($P < .05$). Even in the presence of VPBs, smokers had significantly higher total mortality ($P < .05$) than nonsmokers. A higher incidence of acute coronary deaths in smokers with VPBs compared to nonsmokers with VPBs was also found, but the difference was not statistically significant ($P \sim .08$). Thus, in survivors of myocardial infarction, smoking was associated with a significantly increased mortality, and when complicated by the presence of ventricular premature beats, resulted in an incidence of sudden death greater than the sum of the mortality from sudden death attributed to either risk factor (cigarette smoking or VPB) alone (figure 1). In this study, "lethal interactions" of this nature were found only for smoking, digitalis therapy, cardiac enlargement, and heart failure. These results indicate that cigarette smoking is an




FIGURE 1.—Mortality for men with and without ventricular premature beats (VPB), according to presence or absence of smoking as a risk factor. Bars show percentages of men who died during 3-year period.

SOURCE: Coronary Drug Project Research Group (CV12).

important risk factor in the occurrence of sudden death as well as death from all causes among patients with known CHD.

In a recent article, Mennotti and Puddu (*CV 26*) reported on the 10-year follow-up data of the 1,717 men between the ages of 40 and 59 who were part of the Seven Countries Study. Whereas at five years, no relation could be found between the incidence of CHD and cigarette smoking, the 10-year follow-up data revealed a correlation between cigarette smoking and CHD, with a relative risk of 1.30-1.62 for smokers of greater than 10 cigarettes per day; the relative risk varied with the stringency of criteria for diagnosis of CHD and the degree of smoking. In these two populations, the incidence of CHD was far below that found in the United States, yet the current data suggest that cigarette smoking is a significant risk factor in the development of CHD in Italy.

With regard to the relative risks of the various risk factors, analysis of the 12-year follow-up data from The Peoples Gas Company Study (*CV 31*) revealed that in the cohort of 903 men free of CHD at the initial examination, the risk factor which was associated with the highest 12-year mortality from all causes was cigarette smoking, whether alone or in combination with the other risk factors (table 1). For all "CVR" (cardiovascular-renal), sudden, and CHD deaths there was a trend toward increased mortality in the smokers, but the total number of cases in each category was small.

Carlson and Bottiger (*CV 10*), in the Stockholm Prospective Study, examined 3,168 men and found that, over a 9-year follow-up period, the development of fatal and nonfatal myocardial infarction and other coronary deaths was related to cigarette smoking, elevated serum cholesterol, and elevated serum triglyceride concentrations. The incidence of new events of ischemic heart disease was significantly higher for all smokers ($P < .01$) and for smokers less than 60 years old ($P < .001$) compared to nonsmokers. Smokers aged 35 to 60 had significantly higher serum cholesterol and triglyceride levels, and higher blood pressure readings than nonsmokers.

In terms of CHD morbidity and mortality, cigarette smoking exerts its most potent effect on men under 60. Blacket, et al. (*CV 7*), in a retrospective study of 70 men in New South Wales with CHD, diagnosed between the ages of 28 and 40, demonstrated that 27 percent of this "coronary group" smoked more than 30 cigarettes per day as compared with 11 percent of the controls ($P < .001$). A total of 23 percent of the coronary group smoked more than 40 cigarettes per day ($P < .01$).

TABLE 1.—1958 serum cholesterol, diastolic blood pressure, cigarette smoking status, and 12-year mortality—cohort of 903 men, age 40 to 59 in 1958, free of evidence of organ system disease and ECG abnormalities and followed long-term without systematic intervention. Peoples Gas Company Study, 1958 to 1970.

Serum Cholesterol*	Diastolic Blood Pressure*	Cigarette Smoking*	No. of Men	12-Year Mortality					
				All causes		All CVR diseases	Sudden death	CHD	Stroke
NH	NH	NH	208	15,†	69.9‡	2, 8.1	0, 0.0	2, 8.1	0, 0.0
H	NH	NH	106	11,	80.4	5, 40.2	1, 7.8	3, 21.3	1, 6.6
NH	H	NH	40	3,	72.6	1, 23.8	1, 23.8	1, 23.8	0, 0.0
NH	NH	H	289	38,	135.3	17, 61.3	6, 21.1	13, 46.5	4, 14.7
H	H	NH	34	4,	83.9	1, 20.8	0, 0.0	1, 20.8	0, 0.0
H	NH	H	154	26,	172.2	15, 98.2	7, 44.8	10, 63.5	4, 26.6
NH	H	H	40	8,	170.7	2, 42.1	1, 22.4	1, 22.4	1, 19.7
H	H	H	32	10,	319.3	4, 118.0	1, 26.3	3, 76.4	1, 41.6
1 only high			435	52,	118.0	23, 53.2	8, 18.7	17, 39.4	5, 11.2
Any 2 only high			228	38,	162.8	18, 78.5	8, 36.9	12, 53.5	5, 20.4
Any 2 or all 3 high			260	48,	182.2	22, 84.1	9, 35.3	15, 57.6	6, 22.2
All			903	115,	124.1	47, 51.8	17, 19.3	34, 37.7	11, 11.7

*NH, not high; H, high for serum cholesterol, ≥ 250 mg/dl at entry (1958); for diastolic blood pressure, ≥ 90 mm Hg at entry (1958); and for cigarette smoking, ≥ 10 cigarettes/day at entry (1958).

†Number of deaths.

‡Age-adjusted death rate/1000.

SOURCE: Stamler, J., et al. (CV 21)

The Boston Collaborative Drug Surveillance Program conducted a retrospective study on the prevalence of coffee drinking and cigarette smoking in 276 patients with a diagnosis of acute myocardial infarction (CV 8) and another study on 440 patients with acute MIs (CV 21) comparing these patients with hospitalized controls. On the basis of their data, they suggested that the association between coffee drinking and acute myocardial infarction was stronger than that between cigarette smoking and acute MI. There was observed an almost twofold risk of developing acute MI in noncoffee drinking patients who currently smoked more than 1 pack per day as compared with noncoffee drinking nonsmokers.

Hrubec (CV 20) reviewed data from the veteran twin panel of The National Research Council on patients with angina pectoris diagnosed by questionnaire, and found no significant correlation between coffee consumption and angina, except in the group smoking more than 1½ packs of cigarettes per day ($P < .03$). There was a significant relationship ($P < .05$) between cigarette smoking of any magnitude and the presence of angina.

II. Studies Failing to Demonstrate a Definite Relationship Between Smoking and CHD.

Cotton, et al. (CV 13), in a retrospective study, examined 91 men with a past history of myocardial infarction (4 months to 10 years). Case material was obtained from the files of two referring physicians in England, and controls were chosen from a regional transfusion service. These workers found no increased prevalence of MI among current smokers, but did report a significantly increased prevalence of MI in men with a past history of smoking ($P < .001$). The authors concluded that the lack of significant difference between the patients and controls in their "current" smoking habits ". . . probably reflects the fact that some of the patients gave up smoking or reduced the amount smoked after their coronary attacks."

Durakovic and Saric (CV 15), in a retrospective study of 998 Yugoslavian industrial employees, found no statistically significant correlation between cigarette consumption and the prevalence of angina pectoris.

Malhotra (CV 25) found no difference between current smoking habits in 44 patients with myocardial infarction and 88 hospitalized age- and sex-matched controls in an Indian population. A significant difference was found between cases and nonhospitalized healthy controls ($P = .02$).

Bruschke, et al. (CV 9) reported follow-up from 5 to 9 years after coronary arteriography of 590 consecutive patients with

significant coronary artery obstructive disease, and found a higher mortality rate in smokers compared to nonsmokers, but the difference was not statistically significant.

In New Guinea, Sinnett and Whyte (*CV 29*) reported a very low prevalence of CHD in a rural tribal area. The incidence of other known coronary risk factors was low, but tobacco was smoked by 73 percent and 20 percent of males and females, respectively. However, the authors emphasized that the tobacco leaf is dried and rolled into a bamboo pipe at least 6 inches long "and the smoke is not inhaled."

De Soldati, et al. (*CV 14*) conducted a retrospective study on 66 patients with recurrent myocardial infarction in Buenos Aires, and found a statistically significant correlation between cigarette smoking and recurrent myocardial infarction, but also reported that the nonsmokers had a significantly shorter interval between infarctions than the moderate or heavy smokers.

III. Studies in Women Relating CHD and Smoking.

Most of the studies on the relation between the various risk factors and CHD morbidity and mortality have been done with male populations. Data from large-scale studies have shown a relationship between CHD mortality and smoking in women. In a recent retrospective series of 182 women who died suddenly and unexpectedly, Spain, et al. (*CV 30*) reported a significant correlation between cigarette smoking and sudden death attributed to CHD. In this study, there were 29 CHD-related deaths. There was a significantly greater number of nonsmokers dying of non-CHD causes than of CHD ($P < .01$) (table 2). In addition, 62 percent of the women dying suddenly from CHD smoked greater than 1 pack per day, as opposed to 28 percent of the control group composed of women dying suddenly of other causes ($P < .01$). In the CHD group, there was an inverse relationship between the degree of cigarette consumption and mean age at death, i.e., heavy smokers died of CHD at an earlier age than nonsmokers. The statistical significance of this observation was not recorded. The authors reported that the mean age at the time of sudden death for the entire group of 182 women was 19 years less for heavy smokers compared to nonsmokers. The authors also reported that in their autopsy populations the male:female mortality ratio for sudden death due to CHD was 11:1 for nonsmokers and ~3.8:1 for all smokers.

TABLE 2.—*Cigarette smoking and sudden death from CHD in women*

	Population* Number/percent	CHD sudden death	
		Number/percent	Mean age/range
Smoking habits			
Nonsmoker	81/53	3/10	67/60-75
<20 cigarettes/day	29/19	8/28	55/52-61
≥20 cigarettes/day	43/28	18/62	48/32-58
Total	153/100	29/100	54/32-75

*All deaths exclusive of CHD.
SOURCE: Spain, D. M., et al. (CV 30).

Wink and Hager (CV 35), in a retrospective study of 10 premenopausal women with myocardial infarction, showed that there was a significantly higher incidence of cigarette smoking in this group of women than in postmenopausal women with acute myocardial infarction seen in the same clinic ($P < .001$).

CARBON MONOXIDE (CO)

Introduction

The importance of the effects of low levels of exposure to carbon monoxide on the pathophysiology of CHD is well recognized (1967, 1968, 1969, 1971, 1972, 1973).

It is known that the gas phase of cigarette smoke contains approximately 1 to 5 percent CO and that the concentration of CO increases as the cigarette burns down. Heavy cigarette smokers develop significantly higher COHb concentrations than nonsmokers, reaching levels of 4 to 15 percent. Arterial hypoxemia may develop following CO exposure, and myocardial metabolism may be significantly altered under the conditions of CO exposure, thus limiting myocardial work capacity. A significant decrease in exercise performance in patients with angina has been induced by smoking non-nicotine cigarettes, with increases of carboxyhemoglobin concentrations to approximately 8 percent. Some of the mechanisms responsible for these effects have been reviewed in previous editions of this report.

In experimental studies, CO increases coronary flow and heart rate in normal subjects, increases oxygen debt, and results in myocardial hypoxia. Carbon monoxide has also been implicated by some workers as an etiologic factor in the development of atherosclerotic lesions. Recent studies have contributed significantly to our understanding of the role of CO in the development and pathophysiology of CHD.

Epidemiologic Studies

Wald, et al. (CV 34) found, in cross-sectional analysis of 950 Danish smokers, a correlation between the development of atherosclerotic diseases and carboxyhemoglobin levels. They concluded that COHb levels were better indicators than smoking history of a person's risk of developing ASCVD, including CHD. The heavy smokers had a higher prevalence of ASCVD than the nil and light smokers combined ($P < .001$). Only 8 of the 58 subjects had a past history of myocardial infarction.

Experimental Studies in Man

Aronow, et al. (CV 3) studied the effect of carbon monoxide exposure on myocardial work capacity and angina. Ten men with known CHD and angina, aged 40 to 56, who were exposed to heavy morning freeway traffic in Los Angeles for 90 minutes, performed cardiopulmonary tests in the pre-exposure, immediate post-exposure, and 2-hour post-exposure states. Each man served as his own control by breathing purified compressed air on a similar 90-minute excursion in heavy morning traffic. After breathing the freeway air, all 10 men developed marked increases in expired-air carbon monoxide levels ($P < .001$), and arterial carboxyhemoglobin levels ($P < .001$), both immediately and at 2 hours post-exposure. No differences were found in these parameters when the patients breathed compressed air. The mean exercise performance of the 10 men was significantly impaired in the immediate post-exposure period ($P < .001$) and also at 2 hours post-exposure ($P < .05$). The mean exercise performance was significantly increased from the immediate post-exposure period to the 2-hour post-exposure period ($P < .05$), corresponding to significant decreases in mean expired CO and percent COHb at 2-hours post-exposure. Significant decreases in heart rate ($P < .001$), systolic blood pressure ($P < .01$), and the product of the systolic blood pressure times the heart rate ($P < .001$) developed in relation to the onset of angina in the immediate post-exposure period, and significant decreases were noted at the time of onset of angina at the 2-hour post-exposure periods for heart rate ($P < .001$) and the BP x HR product ($P < .001$). No significant differences in any of these values were found when the patients breathed compressed air. Three of the 10 patients developed ischemic ST segment depression during the ride of carbon monoxide exposure; none of the 10 patients developed such changes during exposure to compressed air. As the BP x HR product is one measure of cardiac work capacity, these results strongly suggest that "less work can be done before the onset of

exercise-induced angina in patients with elevated carboxyhemoglobin levels" (as manifested by diminution of both cardiac work measurements and actual exercise performances). Pollutants other than carbon monoxide may have contributed to the findings in this study.

Using a double-blind design, Aronow and Isbell (*CV 4*) studied the effects of the administration of purified CO and purified compressed air on 10 men with stable angina. All the subjects were nonsmokers. Administration of the compressed air for 2 hours resulted in a significant decrease of venous COHb ($P < .001$) from a mean of 1.07 percent to 0.77 percent and no significant changes in mean exercise time until onset of angina, systolic blood pressure (BP), heart rate (HR), or the BP x HR product. Administration of 50 p.p.m. CO for 2 hours resulted in a significant increase in venous COHb ($P < .001$) from a mean of 1.03 percent to 2.68 percent, and significant decreases in mean exercise time until onset of angina ($P < .001$) systolic blood pressure, heart rate, and the BP x HR product at the time of angina ($P < .001$). It is important to note that the levels of carboxyhemoglobin observed in the groups of patients from these two studies were below those frequently attained by cigarette smokers.

In a related study, Fortuin, et al. (*CV 1, 2, 16*) analyzed mid- and post-exercise ECG changes in "normal" subjects and patients with stable angina before and after exposure to purified CO. In the "normal" subjects, venous COHb concentrations of 5.7 to 7.1 percent were obtained, and exercise performance, as measured by time period of exercise before attaining a specified heart rate, was significantly impaired in both young ($P < .005$) and middle-aged subjects ($P < .01$). Seven of the 26 older "normal" subjects demonstrated some abnormality in their ECGs at some stage in the study, and all of these ECG changes (including ST segment abnormalities and arrhythmias) were exaggerated after CO exposure. In the 10 patients with stable angina, venous COHb levels were raised to 2.9 and 4.5 percent after exposure to 50 and 100 p.p.m. CO, respectively. The mean duration of exercise before the onset of anginal pain was significantly shortened following CO exposure at both dose levels ($P < .005$), ST segment depression was deeper in 5 of 10 patients, and, in general, had an earlier onset and longer duration after CO exposure. In addition, duration of pain was significantly lengthened with exposure levels of 100 p.p.m. ($P < .01$). These studies lend further support to the concept that CO exposure resulting in mild to moderate elevations in COHb concentrations may exacerbate exercise-

induced myocardial ischemia in persons with preexisting clinical or subclinical coronary heart disease.

NICOTINE

Nicotine has been found to increase heart rate, blood pressure, cardiac output, stroke volume, and velocity of myocardial contraction. The cardiovascular effects of nicotine has been summarized in *The Health Consequences of Smoking* (1971). Experimental evidence suggests that these effects are mediated through release of catecholamines from sympathetic ganglia and myocardial chromaffin tissue, and that sympathetic ganglionic blockers inhibit these effects. There may also be neurogenic components to the cardiovascular effects of nicotine, as well as effects on regional coronary perfusion.

Experimental Studies

Bizzi, et al. (CV 6) studied the effects of nicotine on adipose tissue in the rat. Rats were administered nicotine intraperitoneally and subcutaneously, and the responses of plasma free fatty acids (FFA) and in vitro lipolysis were observed. Intraperitoneal nicotine tartrate (NT) caused a transient rise in FFA, and subcutaneous NT in 1 percent carboxymethylcellulose also resulted in a rise in plasma FFA in a dose-response relationship. Administration of the vehicle alone as a control was not performed. Inhibitors of lipolysis (including beta blockers) inhibited this stimulatory effect of nicotine on plasma FFA. Of particular note was that in adrenalectomized rats, NT failed to elevate plasma FFA, but did so when corticosterone was administered before sacrifice.

In vitro, NT added to a medium containing adipose tissue did not result in enhanced release of FFA into the medium, thereby implying an indirect effect of NT through enhanced catecholamine secretion as seen in the in vivo situation. Neither acute nor chronic doses of NT resulted in elevated tissue triglyceride concentrations.

These results confirm previous evidence of nicotine's ability, both acutely and chronically, to elevate plasma FFA, this action being mediated through enhanced catecholamine secretion. The new findings concerning the role of adrenocorticoids are of uncertain significance. The precise role of elevated FFAs in atherogenesis remains to be elucidated.

PERIPHERAL VASCULAR DISEASE

A number of retrospective studies have implicated cigarette smoking as one of the major risk factors in the development and progression of peripheral atherosclerosis, arteriosclerosis obliterans (ASO) and thromboangiitis obliterans (TAO). The patency of peripheral bypass grafts may also be adversely affected by cigarette smoking.

The increased incidence of peripheral vascular disease associated with smoking may be in part due to elevated COHb levels. Experimental data suggest that cigarette smoking may chronically decrease peripheral flow capacity and acutely result in closure of precapillary sphincters, vasoconstriction, and decreased blood flow in human connective tissue in vivo.

Recent prospective and retrospective epidemiologic evidence, as well as experimental data add to our knowledge of the association between cigarette smoking and peripheral vascular disease.

Epidemiologic Studies

The 16-year follow-up data of the Framingham Study (CV 18) revealed strong relationships between cigarette smoking and the development of the three major manifestations of atherosclerosis: CHD, atherosclerotic brain infarction (ABI), and intermittent claudication (figure 2). The only exception to this was the absence of correlation between cigarette smoking and ABI in women, but for CHD deaths, MI, coronary insufficiency, and intermittent claudication, the correlations were as strong in women as in men. It appears that cigarette smoking is one of the major risk factors in the development of intermittent claudication (table 3) (CV 24). A higher total incidence over the 16 years of follow-up and a higher annual incidence of intermittent claudication occurred in smokers than in nonsmokers, the latter statistically significant for all age groups of both sexes. When the other risk factors were controlled for utilizing multivariate analysis, the effect of smoking became even more pronounced.

In a retrospective study of 100 patients with peripheral vascular disease evaluated by peripheral angiography and selective coronary angiography, Tomatis, et al. (CV 32) found that 98 percent of patients with aortoiliac disease, 91 percent of those with femoropopliteal disease, and 93 percent with abdominal aortic aneurysms had a history of smoking.

In a population of nondiabetic smokers with evidence of peripheral vascular disease, Linhart, in Astrup, et al. (CV 5), found




FIGURE 2.—Relative odds of developing atherosclerotic brain infarction (ABI), coronary heart disease (CHD), or intermittent claudication (IC) according to levels of cigarette smoking.

SOURCE: Gordon, T., Kannel, W. B. (CV 18).

TABLE 3.—Standardized multivariate regression coefficients of intermittent claudication for various characteristics

Characteristic	Regression coefficient			Mean (and standard error)
	Age 45-54	Age 55-64	Age 65-74	
Men				
Cigarettes/day	0.362	0.618	0.016	0.451 (0.115)
Glucose intolerance	0.323	0.410	-0.115	0.351 (0.075)
Serum cholesterol	0.303	0.314	-0.103	0.259 (0.092)
Systolic blood pressure	0.105	0.318	-0.412	0.178 (0.107)
Left ventricular hypertrophy (ECG)	0.118	0.016	0.231	0.080 (0.087)
Women				
Cigarettes/day	0.731	0.270	0.462	0.405 (0.126)
Glucose intolerance	-----	0.442	0.232	0.390 (0.101)
Serum cholesterol	0.404	0.286	0.113	0.334 (0.096)
Systolic blood pressure	0.540	0.247	0.420	0.356 (0.136)
Left ventricular hypertrophy (ECG)	0.242	0.223	0.304	0.247 (0.097)

Note: Incidence of intermittent claudication:

	45-54	55-64	65-74
Men	20	45	15
Women	5	21	12

SOURCE: Kannel, W. B., Shurtleff, D. (CV 24).

a significant correlation between frequency of severe intermittent claudication and consumption of greater than 15 cigarettes per day ($P < .01$). There was no difference between these heavy smokers and the other smokers for development of gangrene, nor did development of claudication vary with number of years smoking or total number of lifetime cigarettes consumed. Of his 112 patients, 57 stopped smoking after initial diagnosis and treatment, and subsequent progress of their claudication was retarded with no fresh gangrene developing in any of these subjects. The 55 patients who continued smoking suffered from the same rate of complications as seen before treatment.

Experimental Studies

Goldman, et al. (CV 17) reported that the amount of heat loss from the hands of 10 normal subjects was found to be significantly less on exposure to cold after subjects smoked two cigarettes, when compared to the control state ($P < .05$). Smoking exerted no effect on heat loss from the feet of these subjects. These results corroborate previous reports of the vasoconstrictive effects of smoking on the peripheral vasculature. The differences in heat loss between upper and lower extremities were explained by differences in reactivity to vasomotor stimuli, differences in total blood flow, and greater vasoconstrictor tone in the lower extremity.

CEREBROVASCULAR DISEASE (CVD)

Retrospective studies have revealed a correlation between cigarette smoking and morbidity and mortality from CVD in men. No correlation has been found previously for women.

Acutely, cigarette smoking has been reported to increase cerebral flow and decrease cerebrovascular resistance. The exact role of cigarette smoking in the pathogenesis of CVD is unclear. Recent studies are summarized below:

Epidemiologic Studies

Data from a retrospective series of 598 nonpregnant women, ages 15 to 44, with a diagnosis of stroke (CV 11) revealed that 73 percent of the women with strokes were cigarette smokers as opposed to 43.4 percent of the nonhospitalized control group ($P < .001$) and 60 percent of the hospitalized control group ($P < .001$). The combined effect on thrombotic episodes in this study population of both smoking and oral contraceptives was greater

than that exerted by the use of oral contraceptives alone. It was concluded that cigarette smoking contributes significantly to the development of stroke in women, and enhances the effect of oral contraceptives on the development of thrombotic cerebrovascular events.

Analysis of the 16-year follow-up data of the Framingham study (*CV 23*) revealed an increased risk of ABI attributable to smoking in men, with a sixfold excess risk in male smokers aged 45 to 54. The numbers of cases are still too small to draw any firm conclusions from the data. No correlation between smoking and cerebrovascular events was found in women.

In a prospective study of 3,991 longshoremen followed for 18 years, Paffenbarger (*CV 28*) found no correlation between fatal stroke and cigarette consumption. Included in the study group were 59 cases (from a total of 132) of intracerebral hemorrhage.

CIGARETTE SMOKING AND ASSOCIATED CHD RISK FACTORS

There is conflicting evidence concerning the role of cigarette smoking in the chronic elevation of serum lipids. Results from studies of the acute effects of smoking on blood lipids, including FFA, cholesterol, and triglycerides, have also been contradictory.

A negative correlation between cigarette smoking and mean systolic and diastolic blood pressures has been shown in some studies and not others. This apparent correlation has been explained by the negative association between smoking and relative weight.

Recent contributions in this field are summarized below:

In a retrospective study of 42,804 military men in Belgium (*CV 33*), multiple regression analysis revealed that in all age groups smokers of greater than 20 cigarettes per day had a higher serum cholesterol than nonsmokers. This finding was of statistical significance in age groups 20 to 29 ($P < .01$), 30 to 39 ($P < .001$), and 40 to 49 ($P < .01$); i.e., the years of greatest risk of CHD for the combination of hypercholesterolemia and cigarette smoking (figure 3).

Mundy and Cutforth (*CV 27*) studied 85 patients with myocardial infarctions who had survived their infarctions for at least six months. They found no correlation between "lipid levels" and smoking. Only lipoprotein electrophoretic abnormalities were reported, although the authors stated that serum cholesterols and triglycerides were determined.




FIGURE 3.—Serum cholesterol levels of cigarette smokers and nonsmokers in the Belgian military service.

SOURCE: Van Houte, O., Kesteloot, H. (CV 33).

In a prospective study of 10,000 male Israeli civil service workers, aged 40 and up, Kahn, et al. (CV 22) found a positive relationship between smoking habits at the start of the study and the incidence of hypertension over a 5-year follow-up period (P < .01).

THROMBOSIS

The role of cigarette smoking in promoting thrombosis has not been well defined. Hawkins (CV 19) conducted experiments on platelet functions of 30 healthy men and found that the bi-phasic electrophoretic mobility change of platelets was altered immediately after smoking and returned to normal at 15 minutes.

Whole blood coagulation ($P < .05$), rates of initial clot formation, maximum clot tensile strength ($P < .05$), and clot retraction were all altered in the smokers in the direction of favoring hypercoagulability as compared with nonsmokers, but were not altered further in the smokers either immediately or 15 minutes after smoking one cigarette.

SUMMARY OF RECENT CARDIOVASCULAR FINDINGS

1. Data from recent epidemiologic studies suggest that cigarette smoking acts independently of and in conjunction with certain cardiac arrhythmias to increase the risk of mortality from coronary heart disease in men. Smokers also have a greater probability of dying from CHD at an earlier age than nonsmokers.
2. New epidemiologic data suggest that women who smoke cigarettes have a greater risk of sudden death from CHD than do nonsmoking women.
3. The results of experimental studies demonstrate that the elevated levels of carboxyhemoglobin frequently seen in smokers may result in significantly decreased cardiac work performance and precipitation of ischemic electrocardiographic changes and arrhythmias in patients with clinical and subclinical CHD. Carboxyhemoglobin levels may be of value in determining a person's risk of developing arteriosclerotic cardiovascular disease.
4. Findings from experimental studies confirm that nicotine acts indirectly to cause elevations of plasma FFAs. The relative role of sympathetic versus adrenocortical stimulation of the rise in FFAs remains to be determined.
5. Epidemiologic data reveal strong associations between cigarette smoking and development of peripheral vascular disease.
6. Data from epidemiologic studies support a strong association between atherosclerotic brain infarction and cigarette smoking in premenopausal women and in men of all ages. No association between ABI and smoking has yet been demonstrated in menopausal women.

CARDIOVASCULAR DISEASE REFERENCES

1. ANDERSON, E. W., ANDELMAN, R. F., STRAUCH, J. M., FORTUIN, N. J., KNELSON, J. H. Effect of low-level carbon monoxide exposure on onset and duration of angina pectoris. A study in ten patients with ischemic heart disease. *Annals of Internal Medicine* 79(1) : 46-50, July 1973.

2. ANDERSON, E. W., STRAUCH, J. M., ANDELMAN, R. J., FORTUIN, N. J., KNELSON, J. H. Effects of low level carbon monoxide exposure on human cardiac function. II. Subjects with angina pectoris. Preliminary paper—Environmental Protection Agency with the C. V. Richardson Laboratory. November 24, 1972.
3. ARONOW, W. S., HARRIS, C. H., ISBELL, M. W., ROKAW, S. N., IMPARATO, B. Effect of freeway travel on angina pectoris. *Annals of Internal Medicine* 77(5) : 669-676, November 1972.
4. ARONOW, W. S., ISBELL, M. W. Carbon monoxide effect on exercise-induced angina pectoris. *Annals of Internal Medicine* 79(3) : 392-395, September 1973.
5. ASTRUP, P., et al. Exercise and smoking. IN: Wolf, S. (Editor). *The Artery and the Process of Arteriosclerosis: Measurement and Modification. The second half of the proceedings of an interdisciplinary conference on fundamental data on reactions of vascular tissue in man.* Lindau, West Germany, April 19-25, 1970. New York, Plenum Press, 1972. pp. 119-149.
6. BIZZI, A., TACCONI, M. T., MEDEA, A., GARATTINI, S. Some aspects of the effect of nicotine on plasma FFA and tissue triglycerides. *Pharmacology* 7(4) : 216-224, 1972.
7. BLACKET, R. B., LEELARTHAEPIN, B., PALMER, A. J., WOODHILL, J. M. Coronary heart disease in young men: A study of seventy patients with a critical review of etiological factors. *Australian and New Zealand Journal of Medicine* 3(1) : 39-62, February 1973.
8. BOSTON COLLABORATIVE DRUG SURVEILLANCE PROGRAM. Coffee drinking and acute myocardial infarction. *Lancet* 2(7790) : 1278-1281, December 16, 1972.
9. BRUSHKE, A. V. G., PROUDFIT, W. L., SONES, F. M., JR. Progress study of 590 consecutive nonsurgical cases of coronary disease followed 5-9 years. II. Ventriculographic and other correlations. *Circulation* 47(6) : 1154-1163, June 1973.
10. CARLSON, L. A., BOTTIGER, L. E. Ischaemic heart-disease in relation to fasting values of plasma triglycerides and cholesterol. Stockholm prospective study. *Lancet* 1(7756) : 865-868, April 22, 1972.
11. COLLABORATIVE GROUP FOR THE STUDY OF STROKE IN YOUNG WOMEN. Oral contraception and increased risk of cerebral ischemia or thrombosis. *New England Journal of Medicine* 288(17) : 871-878, April 26, 1973.
12. CORONARY DRUG PROJECT RESEARCH GROUP. Prognostic importance of premature beats following myocardial infarction. Experience in the coronary drug project. *Journal of the American Medical Association* 223(10) : 1116-1124, March 5, 1973.
13. COTTON, S. G., NIXON, J. M., CARPENTER, R. G., EVANS, D. W. Factors discriminating men with coronary heart disease from healthy controls. *British Heart Journal* 34(5) : 458-464, May 1972.
14. DE SOLDATI, L., BARBIERO, O., CHUTLIAN, A., ROMAN, M. I., MATTIAUDA, J., BARREIRO, N. Factores Aterogenicos mas Comunes Observados en 66 Casos de Segundo Infarto de Miocardio. (The more common atherosclerotic factors observed in 66 cases of a second myocardial infarct.) *Presna Medica Argentina* 59(2) : 71-74, January 1972.
15. DURAKOVIC, Z., SARIC, M. Odnos Pusenja Prema Koronarnoj Bolesti u Grupi Aktivnog Stanovnistva. (Smoking habit in relation to coronary

- heart disease in a group of workers.) *Arhiv za Higijenu Rada i Toksikologiju* 22(4) : 309-316, 1971.
16. FORTUIN, N. J., ANDERSON, E. W., STRAUCH, J. M., KNELSON, J. N. Effects of low level carbon monoxide exposure on human cardiac function. I. Normal subjects. Preliminary paper—Environmental Protection Agency and the C. V. Richardson Laboratory. November 24, 1972.
 17. GOLDMAN, R. F., NEWMAN, R. W., WILSON, O. Effects of alcohol, hot drinks, or smoking on hand and foot heat loss. *Acta Physiologica Scandinavica* 87(4) : 498-506, 1973.
 18. GORDON, T., KANNEL, W. B. Predisposition to atherosclerosis in the head, heart, and legs. *Journal of the American Medical Association* 221(7) : 661-666, August 14, 1972.
 19. HAWKINS, R. I. Smoking, platelets, and thrombosis. *Nature* 236(5348) : 450-452, April 28, 1972.
 20. HRUBEC, Z. Coffee drinking and ischaemic heart-disease. (Letter). *Lancet* 1(7802) : 548, March 10, 1973.
 21. JICK, H., MIETTINEN, O. S., NEFF, R. K., SHAPIRO, S., HEINONEN, O. P., SLONE, D. Coffee and myocardial infarction. *New England Journal of Medicine* 289(2) : 63-67, July 12, 1973.
 22. KAHN, H. A., MEDALIE, J. H., NEUFELD, H. N., RISS, E., GOLDBOURT, U. The incidence of hypertension and associated factors: The Israel ischemic heart disease study. *American Heart Journal* 81(2) : 171-182, August 1972.
 23. KANNEL, W. B., GORDON, T., WOLF, P. A., MCNAMARA, P. Hemoglobin and the risk of cerebral infarction: The Framingham study. *Stroke* 3(4) : 409-420, July-August 1972.
 24. KANNEL, W. B., SHURTLEFF, D. The Framingham Study. Cigarettes and the development of intermittent claudication. *Geriatrics* 28(2) : 61-68, February 1973.
 25. MALHOTRA, S. L. In search of causes of ischaemic heart disease. *British Heart Journal* 35(1) : 17-23, January 1973.
 26. MENOTTI, A., PUDDU, V. Epidemiology of coronary heart disease. A 10 years study in two Italian rural population groups. *Acta Cardiologica* 28(1) : 66-88, 1973.
 27. MUNDY, G. R., CUTFORTH, R. H. The relationship between serum lipid abnormalities and other major risk factors in myocardial infarction. *Australian and New Zealand Journal of Medicine* 2(1) : 8-11, 1972.
 28. PAFFENBARGER, R. S., JR. Factors predisposing to fatal stroke in long-shoremen. *Preventive Medicine* 1(4) : 522-528, December 1972.
 29. SINNETT, P. F., WHYTE, H. M. Epidemiological studies in a total highland population, Tukisenta, New Guinea. Cardiovascular disease and relevant clinical, electrocardiographic, radiological and biochemical findings. *Journal of Chronic Diseases* 26(5) : 265-290, May 1973.
 30. SPAIN, D. M., SIEGEL, H., BRADESS, V. A. Women smokers and sudden death. The relationship of cigarette smoking to coronary disease. *Journal of the American Medical Association* 224(7) : 1005-1007, May 14, 1973.
 31. STAMLER, J., BERKSON, D. M., LINDBERG, H. A. Risk factors: Their role in the etiology and pathogenesis of the atherosclerotic diseases. IN: Wissler, R. W. Geer, J. C. (Editors). *The Pathogenesis of Athero-*

- sclerosis. American Association of Pathologists and Bacteriologists, Williams and Wilkins, Baltimore, 1972. pp. 41-119.
32. TOMATIS, L. A., FIERENS, E. E., VERBRUGGE, G. P. Evaluation of surgical risk in peripheral vascular disease by coronary angiography: A series of 100 cases. *Surgery* 71(3) : 429-435, March 1972.
 33. VAN HOUTE, O., KESTELOOT, H. An epidemiological survey of risk factors for ischemic heart disease in 42,804 men. I. Serum cholesterol value. *Acta Cardiologica* 27(5) : 527-564, 1972.
 34. WALD, N., HOWARD, S., SMITH, P. G., KJELDSSEN, K. Association between atherosclerotic diseases and carboxyhaemoglobin levels in tobacco smokers. *British Medical Journal* 1(5856) : 761-765, March 31, 1973.
 35. WINK, K., HAGER, W. Myokardinfarkt bei menstruirenden Frauen. (Myocardial infarction in pre-menopausal women.) *Medizinische Klinik* 67(11) : 364-372, March 17, 1972.

CARDIOVASCULAR DISEASE SUPPLEMENTAL REFERENCES

Part I

(Articles which were reviewed but not discussed in the text, providing data confirming well-established relationships between smoking and cardiovascular disease.)

- 73-0400. ADAMS, J. D., ERICKSON, H. H., STONE, H. L. Myocardial metabolism during exposure to carbon monoxide in the conscious dog. *Journal of Applied Physiology* 34(2) : 238-242, February 1973.
- 72-1061. ALAUPOVIC, P. ET AL. Distribution of arteriosclerosis in various populations. IN : Wolf, S. (Editor). *The Artery and the Process of Arteriosclerosis: Measurement and Modification. The second half of the proceedings of an interdisciplinary conference on fundamental data on reactions of vascular tissue in man. Lindau, West Germany, April 19-25, 1970. New York, Plenum Press, 1972. pp. 21-76.*
- 73-0068. BADRAN, A. M., SOROUR, A. H. A retrospective study of some risk factors in the epidemiology of myocardial infarction in Egypt. *Journal of the Egyptian Public Health Association* 47(4) : 221-231, 1972.
- 73-0656. BARBADO HERNANDEZ, F. J., JADRAQUE, L. M., JIMENEZ, D. L., SIEBERT, M. F., ORTIZ VAZQUEZ, J. El infarto de miocardio juvenil. Revision de nuestra experiencia en la unidad coronaria. (Myocardial infarction among young people. A review of new results in a coronary unit.) *Revista Clinica Espanola* 129(2) : 157-168, April 30, 1973.
- 72-0780. BARBORIAK, J. J., RIMM, A., TRISTANI, F. E., WALKER, J. R., LEFLEY, D., JR. Risk factors in patients undergoing aortocoronary bypass surgery. *Journal of Thoracic and Cardiovascular Surgery* 64(1) : 92-97, July 1972.
- 73-0508. BOLLINGER, A., LICHTLEN, P., KAINDL, F., MANNHEIMER, E. Koinzidenzuntersuchungen bezüglich peripherer Arteriopathie und Koronarsklerose. (Coincidence study on peripheral arteriopathy

- and coronary sclerosis.) Wiener Zeitschrift für Innere Medizin und ihre Grenzgebiete 53(9) : 455-456, 458-462, 1972.
- 73-0071. BÖTTIGER, L. E., CARLSON, L. A. Stockholm Prospective Study II—Nio ars uppföljning. (Stockholm Prospective Study II—Nine year follow-up.) Lakartidningen 69(41) : 4640-4644, October 4, 1972.
- 73-0072. BRÜCKERNOVÁ, O. Kouření u dlouhodobě sledovaných nemocných s ischemickou chorobou dolních končetin. (Smoking in a long-term follow-up study in patients with ischemic disease of the lower extremities.) Vnitřní Lékarství 18(6) : 537-541, 1972.
- 71-0362. CANO, J. P., CATALIN, J., BADRE, R., DUMAS, C., VIALA, A., GUILLERME, R. Détermination de la nicotine par chromatographie en phase gazeuse. II. Application. (Determination of nicotine by chromatography in gaseous phase. II. Applications.) Annales Pharmaceutiques Francaises 28(11) : 633-640, 1970.
- 73-0319. CORELLI, F. Buerger's disease: Cigarette smoker disease may always be cured by medical therapy alone. Uselessness of operative treatment. Journal of Cardiovascular Surgery 14(1) : 28-36, 1973.
- 73-0073. DAMM, G., TRAUMANN, K. J. Risikofaktoren und Arteriosklerosezeichen bei Diabetes mellitus. Rheographische Untersuchungen an 100 Diabetikern. (Risk factors and signs of arteriosclerosis in patients with diabetes mellitus. Rheographic examination of 100 diabetics.) Medizinische Klinik 67(16) : 587-591, April 21, 1972.
- 73-0182. DELAHAYE, J. P., BESSET, C., TOUBOUL, P. Evolution spontanée et pronostic des arteriopathies athéroscléreuses des membres inférieurs. (Spontaneous evolution and prognosis of atherosclerotic arterial diseases of the lower limbs.) Archives des Maladies du Coeur et des Vaisseaux 63(12, Supplement 1) : 64-78, 1970.
- 73-0323. DICK, T. B. S., STONE, M. C. Prevalence of three cardinal risk factors in a random sample of men and in patients with ischaemic heart disease. British Heart Journal 35(4) : 381-385, April 1973.
- 72-0785. DÖRKEN, H. Zur Epidemiologie und Klinik des Herzinfarktes. (Epidemiology and clinical aspects of myocardial infarct.) Internist 13(8) : 310-317, August 1972.
- 73-0513. EKBLÖM, B., HUOT, R. Response to submaximal and maximal exercise at different levels of carboxyhemoglobin. Acta Physiologica Scandinavica 86(4) : 474-482, 1972.
- 73-0325. FINGERLAND, A. Eliminace kouření v prevenci chorob kardiovaskulárních. (Elimination of smoking in the prevention of cardiovascular diseases.) Československé Zdravotnictví 29(8) : 312-320, 1972.
- 73-0342. GREEN, P. W. B., HAWKINS, B. S., IRVINE, W. T., JAMIESON, C. W. An assessment of above- and through-knee amputations. British Journal of Surgery 59(11) : 873-875, November 1972.
- 72-0966. HALL, G. H. Effects of nicotine, carbon monoxide and tobacco smoke on regional blood flow in the cerebral cortex. European Journal of Pharmacology 19(3) : 385-388, September 1972.
- 72-1110. HANSTEEN, V., LORENTSEN, E. Induced hypertension in the treatment of severe ischemia of the foot. Circulation 46(5) : 976-982, November 1972.

- 73-0669. HIGGINS, I. T. T., LOCKSHIN, J. D., GILSON, J. C., COCHRANE, A. L., CAMPBELL, H., WATERS, W. E., FERRIS, B. G., JR., OH, M., HIGGINS, M. W. Coronary disease in Staveley, Derbyshire, with an international comparison with three towns in Marion County, West Virginia. *Journal of Chronic Diseases* 25(10/11) : 567-580, November 1972.
- 73-0839. HILTGEN, M., GUERMONPREZ, J. L., BARRILLON, A., MAURICE, P., LENÈGRE, J. Relations entre le degré des lésions artérielles coronaires, la douleur cardiaque et les facteurs de risque dans 100 cardiopathies ischémiques explorées par cinéangiographie coronaire sélective. (Relationship between the degree of coronary artery lesions, cardiac pain and risk factors in 100 cases of ischaemic heart disease explored by selective coronary cineangiography.) *Coeur et Médecine Interne* 11(3). 521-533, July-September 1972.
- 73-0670. ISACSSON, S.-O. Calf blood flow and smoking. IN: TIBBLIN, G., KEYS, A., WERKÖ, L. (Editors). *Preventive Cardiology. Proceedings of an International Symposium, Skovde, Sweden, August 21, 1971.* New York, John Wiley and Sons, 1972. pp. 171-179.
- 72-0668. JOUVE, A., SOMMER, A., AVIERINOS, C. Épidémiologie de l'athérosclérose coronarienne. Les enquêtes prospectives. (Epidemiology of coronary atherosclerosis. Prospective studies.) *Gazette Médicale de France* 79(13) : 2045, 2046, 2049, 2050, April 7, 1972.
- 72-1112. KARVONEN, M. J. Sydän-ja verisuonitaudit—ehkäisy ja varhainen toteaminen. (Coronary heart disease—prevention and early diagnosis.) *Suomen Lääkärilehti* 27(1) : 76-82, 1972.
- 72-0535. KEYS, A., ARAVANIS, C., BLACKBURN, H., VAN BUCHEM, F. S. P., BUZINA, R., DJORDJEVIC, B. S., FIDANZA, F., KARVONEN, M. J., MENOTTI, A., PUDDU, V., TAYLOR, H. L. Coronary heart disease: Overweight and obesity as risk factors. *Annals of Internal Medicine* 77(1) : 15-27, July 1972.
- 73-0081. KHACHADURIAN, A. K., RUBEIZ, G. A., SAWAYA, J. I. Coronary artery disease in the young. A study of risk factors in 100 consecutive cases in Lebanon. *Lebanese Medical Journal* 25(4) : 253-267, 1972.
- 73-0329. KLENSCH, H., RANFT, C., HARRO, W. Die diagnostische Valenz einiger Risikofaktoren-Tests zur Früherkennung koronarer Erkrankungen. (The diagnostic value of several risk factor tests for early diagnosis of coronary diseases.) *Zeitschrift für Kreislaufforschung* 61(11) : 995-1003, November 1972.
- 73-0331. KYI, K. M., MING, C. K. An epidemiological study of ischaemic heart disease in the Rangoon General Hospital. *Union of Burma Journal of Life Sciences* 3(3) : 337-341, 1970.
- 72-1116. LIGHTFOOT, N. F. Chronic carbon monoxide exposure. *Proceedings of the Royal Society of Medicine* 65(9) : 16-17, September 1972.
- 73-0843. LITTLE, J. M., PETRITSI-JONES, D., ZYLSTRA, P., WILLIAMS, R., KERR, C. A survey of amputations for degenerative vascular disease. *Medical Journal of Australia* 1(7) : 329-334, February 17, 1973.
- 72-0970. MARTELLI, A., PINELLI, G., MARIGNANI, A., RIMONDINI, S. L'Infarto del miocardio al di sotto dei 40 anni. (Studio clinico-elettrocardiografico su 22 casi). (Myocardial infarct in people under forty [Clinico-electrocardiographic study of 22 cases.]) *Acta Gerontologica* 22(3) : 3-18, 1972.

- 72-1296. MIHAIL, A., POPESCU, P., RIZESCU, M., CARUNTU, M., BALAUTA, S. Influenta factorilor de risc si de agravare asupra evolutiei infarctului miocardic acut. (Risk factors that aggravate the evolution of acute myocardial infarction.) *Medicina Interna* 24(3) : 291-303, March 1972.
- 73-0519. NEUFELD, H. N., MEDALIE, J. H., RISS, E., GOLDBOURDT, U. Selected findings of the Israeli ischemic heart disease study. *Geriatrics* 28(2) : 134-136, February 1973.
- 72-0387. NIMALASURIYA, A., WIMALARATNE, K. D. P. A clinico-pathological study of coronary heart disease in Ceylon. *Ceylon Medical Journal* 17(1) : 86-90, June 1971.
- 72-1118. NUSSEL, E., HEHL, F. J. Rauchen und koronare Herzkrankheit. (Smoking and coronary heart disease.) *Rehabilitation, Präventivmedizin, Physikalische Medizin, Sozialmedizin* 25(1/2) : 12-15, 1972.
- 72-1135. OSTAPOWICZ, G. Das sogenannte Raucherbein. (The so-called "smokers' leg".) *Rehabilitation, Präventivmedizin, Physikalische Medizin, Sozialmedizin* 25(1/2) : 20-21, 1972.
- 73-0848. PAULSON, O. B., PARVING, H. H., OLESEN, J., SKINHOJ, E. Influence of carbon monoxide and of hemodilution on cerebral blood flow and blood gases in man. *Journal of Applied Physiology* 35(1) : 111-116, July 1973.
- 73-0162. PERNOT, C. L'infarctus du myocarde en milieu ferroviaire. Aspects médico-professionnels à propos de 100 cas. (Myocardial infarction in railway men. Medico-professional aspects concerning 100 cases.) *Archives des Maladies Professionnelles de Médecine du Travail et de Sécurité Sociale* 33(12) : 611-620, December 1972.
- 73-0087. PREIMATE, E. Y. Rasprostranene nekotorykh faktorov riska i ikh sochetanii pri ishemicheskoy bolezni serdtsa v Rige. (Frequency of some risk factors and their combination in ischemic heart disease in the city of Riga. Epidemiological study.) *Terapevticheskii Arkhiv* 44(9) : 35-38, 1972.
- 73-0089. RICORDEAU, G. Traitement médical de l'athérosclérose oblitérante des membres inférieurs. (Medical treatment of arteriosclerosis obliterans of the lower limbs.) *Revue du Praticien* 22(29) : 4037-4038, 4041-4042, 4045-4046, November 11, 1972.
- 73-0191. RIOS-MOZO, M. Arteriosclerosis femoro-poplitea. Clinica y diagnostico. (Femoro-Popliteal arteriosclerosis. Clinical symptoms and diagnosis.) *Angiologia* 24(5) : 186-195, September-October 1972.
- 72-0792. RISSANEN, V., PYORALA, K., HEINONEN, O. P. Cigarette smoking in relation to coronary and aortic atherosclerosis. *Acta Pathologica et Microbiologica Scandinavica. Section A* 80(4) : 491-500, 1972.
- 73-0677. ROSEN, A. J., DEPALMA, R. G., VICTOR, Y. Risk factors in peripheral atherosclerosis. *Archives of Surgery* 107(2) : 303-308, August 1973.
- 72-0391. SERAFIMOVA, J. Koronarnata bolest i pusenjeto niz iskustvata na kardiološkoto oddelenie na internata. (Coronary disease and smoking—experience of the cardiologic department.) *Godisen Zbornik na Medicinskiot Fakultet vo Skopje* 17 : 709-715, 1971.