first visit to — Tübingen

Choice of interest: apologies

An Approach to Developmental Biology

23 Nov. '70

apologies for only visiting developmental biologists

Molecular biology — what lessons can we carry over to cell biology?

dangers — Hinshelwood.

the "phase" of development (kinds of) in one organism(s), especially convenient experimentally,
— then apply results to other cases.

our approach — divide into different problems,
choose one organism for each one.

what problems?

① classical molecular biology of eucaryotes.
repetitive DNA
nucleus RNA
putting etc.
excretion of proteins
etc.

② membranes: ~~study~~ ~~selective~~

③ mitosis. ~~meiosis~~

④ cell movement — how it moves at all.
— & why it moves in a particular direction.

⑤ cell shape

⑥ & cell contact phenomena

⑦ ~~pro~~ hormones ie chemical signals between cells.

⑧ "positional information"

Role of genetics was very useful in molecular biology

may have an "logic" of situation.

~~pro~~ always extremely useful as a tool when
the right conceptual framework has been found.

but not necessarily always useful.

Kempston . Ampthill Road

Bed. General Hospital

South Lge.

Antipathetic

Scarlet Fever Clinic 4

Name Left when expected

Level of explanation in cell biology

not always necessary to go right down to the molecular level.

eg. existence of enzymes - many not need to know exact molecular mechanism of enzyme action.

but usually useful to go down one level in order to prove a ~~model~~ postulated mechanism.

moreover experimental techniques at the molecular level ~~are~~ very powerful

eg. fingerprinting to show if two proteins are related.

Examples of our work

Sydney Brenner (outline only)

Small soil nematode nematodes (caenorhabditis)

~ 1 mm long - 4 day life cycle - 100 to 200 progeny (eggs, worms.

Grow on - E. coli - defined medium.

diploid $n=6$ - 3 months cool - indefinitely in liquid N_2

Chose it Self-fertilising hermaphrodite. - organism suited to transposon

Summary

genetics - excellent - so related species.

neuroanatomy - very hard work

behaviour - ~~last~~ only described not measured

embryology - not yet started; looks difficult (impenetrable eggs, etc)

Th next article heavily on our work.

Peter Lawrence

(Mary Munro) Wigglesworth.

Michael Locke Rhodnius 3 stages S.

5 larval stages → adult

moult



meal of blood

ripples only on adult

body S

bristle rather sparse.

eye and spine : only for a snapshot.

move sideways, S

move ↓ 25

=

90° 25

" "

ie some "averaging"

Simple model : source, sink, gradient (repeatedly repeated)

concentration – gradient = greater slope – wanders – pseudocentres

90° + diffusion : pattern depends on time.

measuring angles of pattern S



SDVK model : after infinite time.

"pumping" constant. one parameter.

better fit.

later does α depend on : time : double adult

" moulting

: cell "division" : – doubling.

computer simulation.

SLIDES

① Rhodamin - 3 stages

② " - Soda

③ ⟷

④ ↕ equiv.

⑤ ↕ diff.

⑥ g_0 =|||=

⑦ $g_0°$ 

⑧ ~~Eq~~ $g_0°$ Sonk, expr. diff.

⑨ $g_0°$ Sink - relaxed.

---

curr ~~Sink~~ ~~to~~ Runs $g_0°$ ok.

~~Computation~~ Sink here $g_0 180°$ result asymetrical.

here not run asymetrically after ! result.

trying simulations.

Sonic model — here source-sink appears stronger.

Cell motility — mainly work of others
how rapidly developing.

Actin Structures that due to an actin-myosin system.

Hugh Huxley — acellular slime mould.
extracted actin
decorated actin with rabbit heavy meromyosin.
angle, spacing, ATP chars.

Holtzer did it in glycerated cells — chick
+ then decorated. [eg fibroblasts, neuroblasts etc.]

Denis Bray + Dirk Fine

from embryonic brain of chick — decorated with Hugh Huxley

two soluble proteins there in large amounts
— one (55,000) binds colchicine / chemotaxis
— one (45,000) may be actin. — hyper-pinata.

Cytochalasin B Carter (Nature, Jan 1967)

model: stops movement of fibroblasts — quick
— reversible.
tends to flatten cells.
— produces bi (multi) nucleate cells.

Wessells + co-workers (recent PNAS)

stops movement of growth-cone of nerve cells
— confirmed by us etc.

now yes known for — acellular slime moulds
— plants
etc.

claim disorganization.

Thus two aims (1) find (molecular) effect of actin of cytochalasin in the test-tube
(2) natural history of fibrillar molecules in cells.

Reason for meeting: to discuss, and a
to provide discussion held earlier.

our interests

mainly development (nd. nervous system)

also, but less so at the moment, ~~later~~ after d nervous system.

approach

① avoid classical problems.

② tackle problem esp. eg. classical mol. biol. (for ene.)
positional information.
~~chemotaxis~~ swelling gradients
~~Growth~~ cell movement
cell shape
cell division.

& components: cell surface,
membrane receptors & signal lipids
fibrillar elements. permeance

choose "best" option for each sub-problem
(low, or expected dividends).

The perfect problem
- significance
- no long timescales
= techniques available, & easily devised
- interesting results certain
- no one else working.

Our work: our immune system

no-one else working
① neural crest, because invasive.
results - behaviour, mol.
- development ?
② tissue culture } probably explored, but
+ analysis } little - difficult ??

others working
- fibrillar elements. - importance of ~~tools~~ substrates
- cell fusion
- ~~Rous~~ cell surfaces
- BUdR

lead us what answer require

or the visit

no pores in molecular structure unless it helps us

the cellular level

nor always easy to see how deep to probe

~~any~~ can

(although all molecular process are of interest)

embryology of multicellular organisms

many problems - eg. gene control - nucleic acid?
hormones.
cell migration eg. pigment cells.
mitosis; control of.

but also how to determine shape
and specific connections eg. ~~nerves~~ nerves.

neuronal specificity

Sperry; early experiments on adult newts: rotated eyes.
Studied ① behaviour
② histology
Gaze + Jacobson, on developing xenopus.
① early
② later.
Studied by electrodes in tectum + visual stimulation.

found diff. times for the two axes.

ie. not due to function.
postulated two systems of coordinates in eye

[recent results on ipsilateral connections]

How does an eye set up "coordinates"?

① ~~Hoot~~ phase model of Ronald Cohen + Bryan Gaudari.
ie send a constant velocity signal.

② "gradient" models. [historical - got a bad name]
& for example, due to diffusion.

Source + sink - reading concentrations.
Never found how.

is diffusion fast enough? random walk $\frac{L^2}{T}$

values of D. water 20°C. small molecule. 5×10^{-6}

small protein 1×10^{-6}

distances in cell? between cells?

times involved – poh. c few hours.

Suppose distance about 1 mm.

c Sonic Hedgehog. (Spetzai)

Wolpert statements:

typically 50 cells. – broader

(up to 100) in 500μ = $\frac{1}{2}$ mm

Size quoted 20 cells.

Note however that it is the process

many routes
both ie signals
for big distances.

– eg. stripe
colonial aerobe.
(– mushroom growth)

ie. diffusion does mean
an actual diffusion of a marked molecule.

really bilateral alternative – random walk
v. one directional alternative – wave

other explanations

diffusion models (not. work)

– pumping mechanisms.

– all cells known source. sources

Strategy

very unlikely to be able to prove

a model by results from simple manipulation (cut + splice)

because can easily make more complicated models.

Hence valuable because they suggest new types of experiment.

eg. to see what goes from cell to cell

eg. type junctions

(2) are cells polar inside.
microtubules?

(3) can added channels interfere in an assay system.

Model Systems

3D looks too complicated

2D – sheets of cells. tissue culture. one layer of cells.

Drosophila wing – Peter Lawrence

1D System. Sydney. Richard Gibson.

algae – will it be relevant
too early to say.

Finally the, for nerves, other problems

– the reading of the conductance
– transmission to giant axon. (the idea of the addressin)
– synapses, to make right contact.

Harris, Bristles etc.