1/

Rothamsted. 2 March 73.

Our Approach to Developmental Biology

"What's going on in your lab?"

(Sorry, no shirley)

Sydney Brenner: His change from molecular biology of ~~nucleo~~ prokaryotes to development biology.

Plan of Talk: A General Remarks B Examples of our work in this area.

our interest in the nervous system:

how does it work, esp. in man? very difficult problem.

how is it built - one nearer our interests.

general problems of how cells ~~mov~~ divide, move, interact, form tissues, organs and whole organisms.

how to change fields - general remarks.

danger of carryover inappropriate ideas.

Sir Cyril Hinshelwood, from physical chemistry to bacteriology.

excessive ~~exclusive~~ reliance on equations } → "by one bold mathematical assumption
not enough biochemistry + genetics } he managed to ignore the
fundamental problem"

Was not a stupid man

physicists to mol. biology - throw away most previous knowledge.

However this is likely to be less ~~true~~ useful for the move from mol-biol. → cell biol.

What are the lessons of mol. biol.?

① firm theoretical substratum { physics, chemistry, ~~phys~~ physical chemistry

for all biol, mol. biol. is the substratum.
only partly available for higher organisms.

e.g. { v. genetic code – ok.
& { chromosome structure – not yet
(nature of a gene

(as to our level)

② lucky to have a broad theoretical framework, fairly easily
DNA ⇝ RNA ⇝ protein etc.

③ choice of simple systems eg. microorganisms
(in spite of obvious disadvantages)
concentration: eg. T even phages

④ use of modern (often simple) physico-chemical techniques
eg chromatography, ultracentrifuge, radioactive tracers,
electron microscopes, etc.

⑤ the use of genetics as a tool See letter

⑥ the use of cell-free systems: partly for purity (Hoagl otherwise very pure)
partly for ease of addition.

⑦ use of antibiotics as research tools.

General approach S.B. & F.H.C.C.

- reformulate problem and, especially, divide into sub-problems.
  (prob. no easy & unique way to do this)
  for example: can be done in terms of function
  or of structures. (components) (eg.
  omit enumeration: examples later.

- probably wise to accord "classical" problems
  and perhaps "classical" organisms.
  (often chosen long ago, and for reasons which may no longer apply)

How to choose an organism

no general rule: typical questions: how big?
how easy to put electrodes or pipettes into it?
how short a life-cycle?
can it all be done in the lab.?
can one grow (or get) a lot of it?
radioactive labelling?
is biochemistry easy (acellular slime moulds)
how "simple" is it (lower eukaryotes)
is its genetics easy?
worked out? haploid? diploid?

how useful is genetics ? not easy to guess.

may show (or hint at) the overall logic of the problem
eg. Mendel
one gene - one enzyme.

eventually likely to be a powerful tool when.
① the right general conceptual scheme.
② certain technical difficulties overcome
ⓐ "diploid"
ⓑ too many genes.

Overall requirements of the problem

Scientists are human - like to enjoy their work.
typically they would prefer - significance
- no very long investment
- techniques available or easily developed.
- interesting results "certain"
- no one else working ("a few friends")

impossible ! or if it exists, won't stay that way.

5

Examples of work in our laboratory

our interest in the nervous system..

Sub-problem: how do cells know where they are in a tissue?

(colloquium this morning)

(1) insect ~~&~~ epidermis: ~~(1) cells know~~ 2D system. Peter Lawrence

move bits of tissue around & see change in pattern.

conclude: (1) cells "know" where they've come from

(2) " 'forget' their ~~&~~ when they divide

(3) behave as if there was a gradient of

some sort: notion of gradient not yet known

— also, bristle point down the gradient.

no biochemistry yet.

(2) "spacing patterns" eg bristles on the insect cuticle.

1.D work on ~~cyanobacteria~~ anabaena, one of the blue-green algae

Richard Wilcox & Graeme Mitchison. cylindrical cotonic.

chains of cells: heterocysts - fix N_2 - never divide.

vegetative cells - always divide.

can be observed externally & photographed at intervals.

∴ can construct ~~by~~ pedigrees.

~~&~~ can do "cut" chop line not "splice" experiments

Have results (1) influences of neighbouring heterocysts. new septum nearest old one.

Using only one (2) ~~to take out~~ unequal division. — only small fem left.

(3) competition between proheta.

chemistry just starting.

Nervous system : "new" experimental animal.

Sidney Brenner +

Small & soil nematode : caenorhabditis elegans. — nerves, muscle, epithelial cells etc.

small egg, several moults, adults ~ 1mm.

3½ day life cycle brood size 100-200 (∴ 4 day doubling time).

Live on E.coli dish. liquid culture. → diploid (n=6) self-fertilising hermaphrodites. — complications.

occasional males.

marking mutants : point mutagen.

1 week later, ¼ for recessive behavioral mutants, ⅔ show behaviour.

+ morphological mutants

mapping (6 linkage groups). complementation groups.

So far — over 1000 separately arising mutants.

— over 100 complementation groups (= genes)

Thus genetics: very favourable — will eventually prob. be better than drosophila.

paper in preparation.

biochemistry easy to grow say a kilogram

- - - label. (autoradiograph)

not E.coli

Studies just beginning : muscle proteins being studied.

eg. DNA Nuclei complexing ~ x 20 E.coli.

Anatomy & esp. neuro-anatomy

~1000 cells ? 300 nerve cells ~ 96 body muscle cells

no. of cells invariant, also axon branches & connections etc.

has to be done by 1km e/m sections.

very hard work. only part of animal too studied intensively so far.

∴ need for a computer. ~~or~~ 35mm film

assistant analyses data ∴ feeds in (eventually feed out)

expect to have a complete nervous system of adult wild type done with about 2 years.

Mutants: some 6 mers.

Some large changes — no dorsal, but two ventral

⇒ no nerve cords

Some with no changes in that region.

Some with small changes esp. branches fasciculation of few cells.

abnormal

diff. indiv. although same cells, how, branching may be different in diff. individuals.

only a beginning.

no cell division after hatch, except for gonadal.

Embryology : need for study : spend nature. cell-lineage

up to 600 cell stage with light microscope.

e/m : distributions ("shells") : fixing ; now ok..

time taken to hatch: within 6 years or so. (later) study of mutants,

8/

Neurophysiology : electrodes look impossible.
(v difficult even for Ascaris)
but can (prob.) stain for neurotransmitters.

Chemotaxis : to Serra wound . cyclic AMP local neurotr signal.
Thin has advantages + disadvantages - likely to become a
standard animal.

Other lines of work.

① Fibrillar elements Denis Bray (+ Dick Fine)

cytoplasmic ~~Actin~~ actin is embryonic nerve cells. (of chick) 20 i.
also in many other cells other than muscle.
(work of Weissels)
probably also in c myosin (slime moulds)
sliding? cytoplasmic flow.
extension of cell processes?
ruffling membrane: nerve growth cone.

② other fibrillar elements : tubules - microtubules.
~~to~~ neurofilaments.

which slide over which?
how do they get to correct points in cell?

9
=

Membranes

recent advances (review by John Singer)

lipid bilayer : not denaturated protein outside

have globular ones stuck in it + through it.

dynamic picture.

Mark Bretscher : shows that the two major proteins in c

red cell go right through

: believes lipid distribute asymetrical.

Events outside, Serine + ethenolamine, inside

(controversial)

' making ghosts.

Bray studying other proteins in membranes in cells.
by protein diffusion.
Brunner goes to study, nerve surface antigens on surface
of cells transformed by chemical carcinogens
— large variety, etc.

~~Theory~~ Dr (other lines) Acetyl

x-ray studies of chromatin ~ Klug + Roger Kornberg.

: biochemistry : Marcus Noll. Sept.

Theory of nervous system : David Marr.

Conclusion.

~~Electron only a small fraction~~

General Remarks

can only tackle a small part of the field.
advantage of having varied lines of attack.

- one new organism nematodes.
- one "new" problem parasitic nematodes.
- the usual population problems eg. fibrillar elements
  cell membranes.
  chromosome structure.
- one "difficult" technique : John Gordon injection into oocytes.
- theory : David Brown (René Thom)

but first a little reading & then how a long way to go

Classical methods — [illegible] — [illegible]
— [illegible]

John Snow & the pump:

Nomenclature
Prokaryotic & eukaryotic. < spans partition.
Fibrillar elements; cell movement.
Cell membranes:
Chromosome structure: — x-rays; — coincident problem.
— chemical composition
Meyer: David Baum: see Henry.