The chemical formula of DNA is ~~well~~ now well established. It ~~consisted~~ is a long fibrous molecule containing purine and pyrimidine nucleotides - bases linked Together ~~by~~ 3-5 phospho-diester linkages at ~~the base~~ 5 ~~positions~~. The back bone of the molecule thus consists of a regular alternation of sugar and phosphate groups as shown in Figure 1. Four different types of nucleotide bases are possible, and it has as is known the sequence of bases is irregular. Two of the possible bases, adenine & guanine - are purines, and the other two - thymine and cytosine - are pyrimidines.

~~Our structure~~ Previous consideration of the structure of DNA Since the chemical formula of DNA consisted of one very long polynucleotide chain, it was generally assumed that its structure could also consist stereochemical of one chain. ~~Crystallographic evidence~~ are doubtfully conclusive however make it ~~most likely that~~ suggested that the ~~crystallographic~~ structure must standard consist of one that one chain. ~~We that~~ We ~~have also~~ postulated a ~~structure~~ of our structure therefore consists of not one but two polynucleotide chains both coiled round a common fibre axis as is shown diagrammatically in Figure 2. The two chains are held together by hydrogen bonds between the bases.

~~There~~ was only one chain in ~~the chemical formula~~

It has often been assumed that since the chemical formula ~~has usually~~ there would only be one in the ~~unit and only one chain has been~~ ~~This would also be true for the~~ However the

structural unit. ~~The X-ray evidence and the density~~ The density,

taken with the X-ray evidence, 4,5 suggests ~~indicates~~ very strongly that ~~the structural unit consists of more than one chain~~ there ~~must be more than one~~ polynucleotide chains are probably two.

We should emphasize that while the sequence of bases is indeterminable, the necessity for specific pairing demands a complementary relationship between the sequence of bases on the opposite chains. ~~That is~~, If the sequence on one chain is given then the sequence on the other is automatically determined. ~~The one chain is, as it were, the complement of the other~~. Our model, consists, in effect of a pair of chains, either of which can be regarded as the complement of each other. This feature appears to us most important as can be seen by rephrasing the previous sentence to read, our model consists of a pair of templates, either of which we can be regarded as the negative of each other. Thus, if a single chain could acty as a template (or mould) for the formation of its complement, the understood mechanism of DNA self duplication would in principle be solved.

The idea that a single chain could act as a template for the formation of its complement seems reasonable. Our model suggests that this could be done most simply if the single chain (on the relevant portion of it takes up the helical configuration of our structure.

Under out the above scheme, ~~of autoreproduction~~ the sequence of the pairs of bases will ~~have been~~ be duplicated exactly. ~~That is as there~~ ~~the necessity for spontaneous pairing~~. If, however, a bases, occurs in one of the less likely tautomeric forms, at the moment when the complementary chain is formed a mistake in copying occurs will result. For example adenine will normally pair with thymine, but if there is a tautomeric shift of one of its hydrogen atoms it can pair with cytosine. The next time duplication occurs the adenine would pair with thymine, but the cytosine will guanine. And we do not know the exact energy difference between the tautomeric form we consider as yet precise with what frequency mistakes might occur. If the energy difference were not more than 20 Kcal, then a mistake would occur one in 10^9 of one pair of bases, and if we assume there are 1000 pairs of bases in a gene duplication, and that have a frequency comparable to the natural of spontaneous mutations

we feel.

of our structure. ~~We begin~~. Under this hypothesis we would imagine that ~~at~~ at this stage in the life of cell free nucleotides (~~strictly~~ polynucleotide precursors) become available are join up by hydrogen bonds to one of the bases on the ~~the~~ already formed chain. We would postulate that the polymerization of the ~~next~~ nucleotides to form a new chain is ~~only~~ possible if the resulting chain can form the proposed structure. This is plausible ...

That is to satisfy the requirement of specificity

Naturally for this process to ~~work~~ be important, at some stage in the cell division, the two complementary strands must separate from themselves, that is, they must untwine themselves. As the ~~turns~~ of the turn of the helix occurs every 34 A, a very large number ~~number~~ of entanglings not occur (in fact ~~is~~ 1500 turn for ~~every~~ per molecular weight of 10^6]. The postulation of such a large amount of uncoiling, however, does not seem to us an unsurmountable objection to our ~~present~~ proposal of autoreproducting, ~~but~~ especially in view of the extensive coiling and uncoiling of chromosomes which occurs during mitosis. While this microscopic coiling is at a scale many fold larger than our molecular unit, it does but seem unlikely that this coiling ~~could~~ ~~can be~~ is based on a ~~the~~ process at the molecular level.

It is of interest to consider what would happen if a base occurred
~~should happen to~~ is one of its less likely tautomeric forms at
the moment when the complementary chain is being formed. ~~For example~~
While adenine, for example, will normally pair with thymine, if there
is a tautomeric shift of one of its hydrogen atoms it can pair
with cytosine. The next time pairing occurs, the adenine
(having resumed its more usual tautomeric form) will pair with
thymine, but the cytosine will pair with guanine, and thus
a mistake will have occurred in the copying process. It seems
may
possible that this might be the cause explanation of
spontaneous mutation. ~~rate.~~ The difference in energy
between tautomers under physiological conditions is not known, but
biological
making plausible assumptions a difference of about 10 kilocalories
would be necessary to obtain the right order of magnitude for the
observed rate. ~~of spontaneous mutation~~.

It is also possible that one chain of a pair, perhaps under the strain of twisting, occasionally breaks. The polynucleotide chain remaining intact could then relax the accumulated twist by rotation about single bonds and following this, the broken ends, being still in close proximity, might rejoin. At present this is not possible to prove nor disprove, but the fact that it would explain the equilibrium between the open circular and supercoiled forms of DNA is at least a clue to this phenomenon. If this process occurs often, the breaks must be very few, since otherwise the DNA would lose its biological activity.

Legends to Figures

1. The chemical formula of ~~DNA, shown diagrammatically~~ a single chain of DNA.

2. This figure is purely diagrammatic. The two ribbons symbolize the two phosphate-sugar chains, and the horizontal rods the pairs of bases holding the chains together. The vertical line marks the fibre axis

3. The chemical formula of a pair of DNA chains. The hydrogen bonding is ~~shown~~ symbolized by dotted lines.

4.