

Reprinted from Abstracts, 7th International Congress  
of Microbiology, Stockholm, 1958, pp. 58-60.

Extranuclear transmission of the F compatibility factor in *E. coli*. Joshua Lederberg, Departments of Genetics and Medical Genetics, University of Wisconsin, Madison 6, Wisconsin, U.S.A.

The mating types of *E. coli* include  $F^-$  females and  $F^+$  males, the latter being determined by the presence of a readily contagious factor "F". A number of lines of evidence converge to suggest that this is inherited outside the chromosomal system of linked markers which encompasses most of the known heredity of *E. coli*: 1) in crosses of  $F^+ \times F^-$ , virtually all of the progeny are  $F^+$  while other markers segregate in characteristic ratios. 2) When an  $F^-$  culture is seeded with a few  $F^+$  cells, the  $F^+$  character spreads through the entire population, indicating that it multiplies more rapidly than the other genes. 3) After pairing with an  $F^+$  cell, an  $F^-$  cell gives rise to a pure  $F^+$  clone; in comparable matings of  $F^- \times Hfr$  (high frequency of recombination with respect to other markers) the exconjugant  $F^-$  clone invariably segregates a mixture of unaltered parental and recombinant types. 4)  $F^+$  is transferred more rapidly and efficiently than any other known markers, and in  $F^+ \times F^-$  crosses, shows no linkage to any other marker. 5) As described by Y. Hirota, the  $F^+$  factor can be efficiently removed from a population of  $F^+$  cells by treating them with acridine dyes, especially acridine orange. These findings suggest that the  $F^+$  cell carries an extranuclear factor, i.e., a plasmid, which is rapidly and efficiently transferred to  $F^-$  cells during brief contact. In many respects, the inheritance of F is analogous to that of the factor which governs the trait colicinogeny as analysed by Fredericq. The mating behavior of Hfr mutants, the instability of

some of these to give infective  $F^+$  reversions and the segregation of the Hfr in various linkage relationships in Hfr x  $F^-$  crosses suggest that the Hfr mating types represent the fixation of the F factor at specific chromosomal loci.

The dearth of recombinants given by  $F^+$  compared to Hfr has provoked the hypothesis that  $F^+$  is intrinsically sterile, and that its recombinational fertility depends on spontaneous mutation to Hfr. Such mutants undoubtedly occur, but it is difficult to assess their role in  $F^+$  fertility without an accurate measure of the mutation rate. The regularity with which  $F^+$  x  $F^-$  crosses give  $F^+$  progeny argues against the necessary role of stable, non-infective Hfr mutants similar to the type Hfr<sub>Cavalli</sub>. Control experiments have shown that the regularity of  $F^+$  progeny is not due to secondary reinfection. It is still possible that in addition to occasional Hfr mutations, the more fertile cells in an  $F^+$  culture have experienced more transient changes in the quality or position of the F agent they carry.