U.W.
MICROBIAL
GENETICS

Hfr
3200 $\frac{F_{13}}{}$ × W3086 W

26

12/IV : 1959

REF:

Method

Overnight culture of W3200 5 ml/ pen + 0.1 ml W3086

Incubate overnight (20 hrs)

purify it on Blacsm-agar.

Replica plate it on MGal seeded W2979 on it.

Result & conclusion : F'_{13} is obtained from Hfr_{13} with the method described above. F' gives Hi for Gal; Medium for Try + A_1; Low for G_1 + TLB_1



on MGal

Comparisons on infectivity of F'+ and corresponding Hfr.

16/4 1959

REF: cf. p. 33. infectivity.

· cultural age : Overnight culture (Penassay)

· Strains: Hfr is isolated by purification of stab culture.
W4321 (Hfr_4 M^-) W3208 (Hfr_8 M^-)

· Experimental method:

Hfr or F'+ 1 ml + 3086 0.2 ml + Penassay 5 ml.

↓ Incubate it overnight.

↓ Purify it on MMal Sm agar.

↓ Test sex-compatibility

Result:

	Hfr / total.	% converted into F'	infectivity
(Hfr) W4321 × 3086	0/24	0	−
W4518 (W6 F_4) × 3086	22/22	100.	+
W3208 Hfr_8 × 3086	18/22 (not Hfr: F+?)	82 (usual F+?)	see back page.
W4520 × 3086	20/25	80	+

Thus non-Hfr of W4321

Conclusion : Original stab culture contains 2 kinds of Hfr strain, infective and non-infective.
Infective Hfr may occurred by mutation of non-infective Hfr × F' is simply splited from chromosome.





Mac.
K2985
A1

K 2979
m Gal.

I should use Gal marker, because no asymtrophy.
try again.

3086 F_8 —× W6 ●

F_8 —× F_1 Is it co-exist or exclud ? P.29
[with each others. P. 48
REF:

16/IV ; 1959.

3086 F_8 : 1ml : F_1 0.1ml : 5ml penassay.
W6
↓
Incubate it for 24 hrs at 37°C.
↓
Purify it on D Mal
↓
Test Mal+ colonies on sex. in cross times 2979. on MGal.

3086 F_8 —× W6



test infectivity of F, and F_8 to F- at 7 (P29)
Does this clone still contain F_1 ?

3086 F_8

× 2979
on MGal

sft/total testes : 1/32 × 100 = 3.12

Conclusion : F_8 is infective to W6 F^+.

objection : W6 —spontaneous loses of F→ W6 F^- → W6 F_8
3086 F_8

W6(F1 F8) ⟶× 3086

23/IJ ; 1959 REF:

Purpose of this experiment: Does W6F1F8 still retains F1 or not?
cf. P.28.

		W6F1F8				
Control 1.	or	W6F1	——×	W3086		
Control 2.	or	W6F8				
		1 ml	+	0.1 ml	+ 5 ml	plasseg

↓

incubate overnight.

↓

Purify it on BLac 5m.

↓

Cross-brush it ⊥ mm × 2979 on M Gal.

		F8/total (%)	F1/total (%)
Experiment:	W6 F1 F8 ×3086	6/30 (20)	0/30 (0)
Control:	W6 F1 (W6) ×3086	0/34 (0)	17/34 (50)
	W6 F8 ×3086	30/34 (88)	0/34 (0)

Conclusion: W6 F1,8 (obtained by infection of F8 to F1+ : W3086F8 × W6)
does not contain F1, or not segregate in the time of infection.
F8 may exclude F1 after infection. F8 > F1
stronger than.

F donor : W6 F_1 , W6 F_8 , W6 $F_{8,i}$



1/10 5/11 10/11 ~~[illegible]~~ ×3828 on Mlec

3/10 6/11 10/12 ×3828 on Mlec

2/10 6/12 10/11 ×3828 on Mlec.

F recipient : 3086.

		%
$F_{1,8}$	6/30	20
F_1	17/34	50
F_8	20/34	88.

U.W.
MICROBIAL
GENETICS

Direct demonstration of true recombination. Not syntrophy!

30

Check syntrophy between Gal⁻ and M.⁻ A and M.

20/IV ; 1959

REF:

1. Mix W6 F_8 1 ml + W2979 1 ml + 5 ml pen. Ca. 10^8 cells/ml.
   ↓
   incubate 1 hours at 37°C.
   ↓
   dilute it into 10^{-1}, 10^{-2}, 10^{-3}, 10^{-4}, 10^{-5}, 10^{-6}.
   and spread it on MGal and BGal.
   ↓
   Incubate 40 hrs and see what happens.

See : back page.

Degree of dilution	10^{-1}	10^{-2}	10^{-3}	10^{-4}	10^{-5}	10^{-6}	survivor cont. 10^{-6}
# of colonies	too many	too many	too many	150	32	6	17

2. pick cells from blackspot and ~~streak~~ streak on BGal ~~is it Gal+ or Gal⁻~~, and test nutritional requirement on ~~MGal (Replica plate from BGal to MGal)~~ replica plate aux/prot.

		isolated on	Test nutritional requirement:	% of x⁺
W6 F_8 × 2979	on M Gal	BGal	almost Gal⁺ tested Gal⁺	0/22	0.0
W6 F_8 × 2985 A^-	on MLac	Bgal.	All gal⁺	4/21	16.0
W1994 F_8 × 2979	on MGal	BGal.	9/3 gal⁺. tested Gal⁺	2/23	8.0

young culture. 4 Spot it on MLac SM B_1 seeded 3928
old culture

SM destruct male capacity of $^{S^S}_{\lambda}F'$ male but not S^r F' male.



Conclusion : ① This phenomenon is not syntrophy but true recombination.

② F8 × F⁻ S^R (recombination process) is inhibitted by SM-treatment. (It may be interpreted by incorporation of S^S loci to the F-S^{SM})

③ F4 × F⁻ S^R is not inhibitted by SM.
This phenomenon may be interpreted by incorporation of S^S to F^{S^R}.

W6 F_4 x W2985



W6 F_8 x 2979



Test of the syntrophy.



W1394 F_8 x 2979



Conclusion:

almost of three Gal+ colonies are

X^+.

Therefore, this phenomena are not syntrophy but recombination.

Genetics.

Preliminary experiment for exclusion of F particles.

U.W. MICROBIAL GENETICS

31

W6 F_4 $\xrightarrow{F_8}$ × 3086 F_8

W6 F_8 $\xrightarrow{F_8}$ × 3086 F_4

16/IV. 1959 REF: cf. p 42 ~ p 50.

Ratio of mix.

Donor (W6 F') 1 ml + Recipient 0.1 ml + Penassay. 5 ml.

↓

Incubate overnight at 37°C

↓

Purify it on B Hard SM.

↓

Test Mal⁻ colonies for sex-compatibility.

The method used in this experiment is not suitable to know the difference between two different F'+. Use spot method for this purpose. (See back page.)





test each H1 markers. Retest H1 markers: See back page

conclusion : F_4 is stronger than F_8. F_4 seems expel F_8 after infection.

But, is it chromosomal level or cellular level? Does phenotypic F_8 (F_8) still contain F_8 or not? (See Back page.)

Next step : $F_{4,8}$ × F^- and see segregation of F_4 and F_8.

Fertility? patterns of F_4, F_8, and $F_4 \times F_8$, $F_8 \times F_4$ obtained by ifation. of F_4' to other F^+





F_4 F_4 F_4 F_4 F_8 F_4 F^-

$F_4 \times F_8$ control.

$F_8^{\sigma} \times F_4$

Phenotype F_4 F_4 F_4 F_4 F_8 F_4 phenotype

$F_4 \rightarrow F_8$ control

$F_8 \rightarrow F_4$.

Isolation of F_2 from Hfr$_2$.

4068 Hfr$_2$ ⟶ × 3086. F⁻ M Mal S^R

16/IV. 1959. REF:

4068 : From Gah. (15/IV) Exponentially growing. it is grown into pm. for overnight

3086 : old. broth

Ratio of mix.

4068 /ml : 3088 0.1 ml : Penassay 5 ml.

↓

Incubate then at 37°C. for overnight.

↓

Replica plated on MLac (×3828) or MLac B; SM. (1394).

↓

Pick High recombination coloring and test Hfr quality (×1394) on MLac B; SM



Replica plated on MLac seeded ×3828

on MLac Sm.

Treatment of W1922 F8 by AO.

22/IV. ; 1959 REF:

Time of treatment : Ca. 40 hrs. at 37°C. Conc. of AO 30 γ/ml.
5 ml passage.

Purpose : W1922 is Hfr. Has W1922 F8 Hfr, or not. If Hfr, is in W1922 F8
it will gives Hfr, after treatment. Tester : W2979 or M Gal.

Result:

	AO treated W1922 F8	Untreated W1922 F8.
+/total	27/27	0/23
% of +	90	0

Conclusion :
W1922 F8 does not give arised Hfr, after AO-treatment. Only gives F⁻.



U.W.
MICROBIAL
GENETICS

36

FR. from F8 3086.

28/N — 1957

REF: cf. p35
P37 51 49

	1	2	3	4	5	6	7	8	9	10
		U.V.	10 sec.		Initial number of cell /plate : ca. 10^7–10^8 cells/pl					
					cultural age : overnight.		seeded on B Gal.			
							strain : W2979			
							med : on M Gal.			
	plate #	# of colonies/plate		# of Rec+ colonies.			% colony appeared.			
	1									
	2									
	3									
	4									
	5									
	6	19		18			1 ↙	pick and retest the leaks of		
	7	26		26			0	fertility.		
	8	36		36			0	and test immunity.		
	9	31		31			0			
	10	90		90			0			
	11	35		35			0			
	12	53		53			0			
	13	37		37			0			
	14	37		37			0			
	15	49		49			0			
	16	63		63			0			
	17	53		53			0			
	18	29		29			0			
	19	25		25			0			
	20	28		28			0			
	Σ.			610			1			

Interrupted mating experiment of W6F8.

18/IV ; 1959. REF: H is low mafer. bac.

F⁻ : W4352 F⁻ Try P M Gal₆ Lac₅₇ U_6^R Lp^+ Make S^R mutant.
Hfr : W4520 Fg H.
(W6F8)

Method.

Selective marker:	Medium	Purpose
Try	M Gal conc + Meth + Prol	Try incorporation.
Try & Prol	M Gal + Meth.	2 Which is earlier Try or Prol.
Try Lac.	M Lac + Meth + Prol	Which is earlier Try or Lac.
"	" + Meth + Prol + Trp	1 Lac incorporation
Try & Gal	M Gal + Meth + Prol	which is top. Try or Gal.
Gal	" " " + Trp	Gal.

Method:
1. Purity W4352 on Blac agar. (It was very small colonies).
2. Inoculate purified colony into peptone and incubate it at 37°C. for overnight.
3. Transfer it into peptone broth (10ml) and inoculate W4520 into 10ml peptone and shake it for 4hrs. (11:00 — 2:45)
4. Mix them (1:1) chilled take a sample, and dilute it into H_2O. ($\times 10^{-3}$, & 10^{-5} dil: ca 10^6 & 10^8 0time 0time viable/ml)
5. Spread over each selective medium at each time.

Selective Medium	Time / Selective marker	2:45 0 min. after mix. (# of cells ×100.)	2:50 5	2:55 10	3:00 15	3:05 20	3:10 25	actual time. min.
Gal-M-P	Try	0	0	0	0	0	0	
Gal-M-P	Try-Gal	0	0	0	0	0	1	
Lac-M-P-Try	Lac	0	0	0	0	0	0	
Gal-M-P	Try-Lac	0	0	0	0	0	0	
Gal-M-P	Gal.	3 ($\times 10^2$)	1	?	5	9	0	
BLac +	Survival Count							# of cells used in Exp one ($\times 10^2$).
F_8	# of 4520 a	0	0	0	0	0	0	
	# b	0	0	2	0	1	0	
F⁻	# of 4352 a	7	12	33	15	12	7	
	# b	12	24	38	18	27	11	

Method of dilution.

0.1/10 × 0.1/10 : 0.1 /plate : 5 min —— 25 min. (selective agar)

0.1/10 : 0.1 /plate : 0 min. only. (selective agar)

0.1/10 × 0.1/10ml × 0.1/10ml : 0.1ml : 0 min —— 25 min (EMB Lac)

U.W.
MICROBIAL
GENETICS

40

Crude timing experiment of for chromosome transfers from F'+ to F−
markers observed : Lac.

24/10 ; 1954.

REF:

Cultural age : ca. 5 hr. on rotator.

cell number : Ca. 10^8 cells/ml.

Phenotype.

selective media : M Lac Sm.

Strain : W6 F_8 ♀ W3828.

dilution		
10^{-3}	10^5 : 0.1ml/10ml	: 0.1ml/plate
10^{-4}	10^8 : 0.1ml/10ml	: 1ml/10ml : 0.1

Time (min) / dilution	0	0	10	20	30	40	50		120
Ca. 10^5/plate 10^{-3}	13	1	10	70	56	81	80		172.
Ca. 10^3/plate 10^{-4}	0	0	0	1	4	5	7		30

0.01 % # of recombinants.
of recombinations.

9
8
7
6
5
4
3
0.02
1

0 10 20 30 40 50 min

U.W.
MICROBIAL
GENETICS

41

W6 F8 × W3828.

Is there 2 states in W6 F8 ?

23/10. 1959 REF:

W6 F8 × W3828. on MLac Sm, & MLac.
M⁻ F8 Lac⁻ S^R.

Hypothesis: Interpretation for low fertility of W6F8 on Sm-media.

: Almost of the F8 cells are in state I (cytoplasmic state), and not state (II).
Only state II can donate chromosome even in the presence of Sm.

: If it is true, all of W6F8 colonies show Hi on MLac, and only (clone) four W6F8 shows Hi fertility in the cross on MLac Sm. (W6F8 only shows the state II phenotype on (MLac) Sm-media) and almost of the W6F8 colonies are low fertile.

Method:

1) Make W6 F8 colonies on BLac agar. (10 plates).

2) Replica plate it on MLac and MLac Sm seeded W3828 on it,

3) Incubate it for 30 hrs.

plate #	# of colonies	Hi # of Fertile colonies on MLac	on MLac Sm.
1	135	135	135
2	112	112	112
3	130	130	~~129~~ 129(1)
4	127	127	127
5	58	~~57~~ 58(1)	58(1)

Result.
apparently, fertility of F8 W6 (on MLac Sm) is much low than MLac. But there is almost no fluctuation among replicated colonies.
all State I ?

F⁻ or F^R =

Conclusion:
This phenomenon (inhibition of recombination formation by Sm.) may be interpreted by the incorporation of S^S segment to S^R F⁻.
(from Hfr parent)



one example.
Mlac
pickad

2nd trial of infection of F_4 — × F_8.

23/4, 1959. REF: Cf. p 31. Cf. 50

• Method : (W6) (W3086)

Mix F_4 1ml : F_8 of 1ml : 5ml phoassy. Control F_8 only.

(W6 F_4 W3086 F_8 is purified on Olac. before using.)

↓ Incubate it overnight.

↓ Purify it on B Mal Sm.

↓ Mal⁻ Sm^R colonies are into 1ml phoassy. tubes. and let it grow for 5 hrs at 37°C

↓ Spot them on (W3086, W2985, W2979) m. Oal.

(Fm) (A_1) (Oal)

and compair Hi recombination pattern.

• Strain : W6 F_4, W6 F_8, W3086 F_4, W3086 F_8 all of them are purified before using in this experiment.

• Result :

Experiment	HI for A_1 only (F_4)	HI for Gal_2 only (3086 F_8)	HI for both (double)
W6 F_4 × W3086 F_8	1 #28 (F_8 may be expelled by F_4)	38	5 1 2 15 31 34
Control W3086 F_8 only	0	40	0

Double F strains. (F_4 + F_8 in W3486)

• Conclusion :

1. F_4 is infectable to F_8^+ cell, and sometimes expel F_8 from the host cell and sometime make double F strain.

2. These fertility ▶ may depends on the segregation of two types F^+. (∵ If A_1 becomes higher than parent Gal will became lower or vice versa, not both.) See segregation of F characters by spot test and replica plating method.

[redacted]

efficient method : application of Replica plating method.

W6 F_4 × 3086 F_8. Use Xyl to detect F_4

Seed on B Mal Sm.

Replica plate on MXyl seeded W2979.

(F_4 is Hi for Xyl but F_8 is very low for Xyl, therefore it is easily differentiate between F_4 and F_8.)

on MXyl W2979

Take HI colony, on Xyl transfer and see segregation of the marker by replica plating in same way.

×2979 (Gal⁻) 2985 (A_1) on MGal

Control W3086 F_8.



× W2985 ($A_1^- F^-$) on MGal



× 2979 ($Gal^- X^+$) on MGal.

infection. Look for double F.

W6 F_4 — × W3086 F_8

× W2985 ($A_1^- F^-$) on MGal



× 2979 ($Gal_2^- X^+$) on MGal

Test No. 28, ●. No. 29. and No 31. No 1.
on. Segregation, or [redacted] H1 for both.

2nd trial of infection of F_8 to F_4^+

$W6\ F_8$ —× $W3086\ F_4^+$.

23/IV 1959

REF: (cf. p 31.) $F_4 \rightarrow F_8$

Method: is completely same as $W6\ F_4$ —× $W3086\ F_8$.

Purpose: Is $W3086\ F_4^+$ accessible to the infection of F_8?

Result:

			H_1 for A_1 only (F_4 type)	H_1 for Bal_2 only (F_8 type)	H_1 for both (double F)
	$W6\ F_8$ —× $3086\ F_4$		35/35 (100)	0/35	0/35
Control	$3086\ F_4$ only.		35/35 (100)	0/35	0/35

All the colonies of 5 tested

Conclusion: [$3086\ F_4$] does not express F_8 character after mixed culture with $W6\ F_8$.

This may be explained as ① lack of infection from F_8 to F_4.

or ② it is infectable to F_4 but recessive for phenotypic expression.

or ③ it is infectable to F_4 but is expelled after infection by exclusion between F_4 and F_8. F_8 is weak to the competition than that of F_4.

U.W.
MICROBIAL
GENETICS

46

Test for 2 states of fertility in $F_8^+ f_3$

W4525 (W3644 F_8^+)

1/M : 1957.

REF: ~~FLB, Hel, SR f_3.~~ Cfp 55, p 70, p 23

W4525.

① Purify $F_8^+ f_3$: (W3644 F_8). on B Lac.

② Inoculate it into penassay and let it grow. for overnight.

③ Make colonies on Blac. and replica plate it on M Gal plates ~~W3644 F M Gal~~ W2979 ~~W3644~~ or W6 F−.

Purpose :

Does $F_8^+ f_3$ shows two states on Gal or Mlac in ♂ or not ?

(only M or both or only Gal ?)



Result :

only one state of fertility was observed. (see below) in both cross. (× M−, × Gal_2 −)

Conclusion : F_8 does not fixed on Rfd_3 locus. Only attach at the Xyl locus

(∵ Lo. for Meth (Rfd make hi). ∴ Hi for Gal. and also only one state was observed.)

This means F_8 does not split off as easily as F_1.



W4525



# of colonies 220

× 2979 on M Gal

# of colonies 220

× W6 F− on Mlac

Minimal cell number for infection of F' to F⁻

5/V ; 1959 REF:

1. Overnight culture (ponassy culture) of W6 F8 and W3086 are inoculated into ponassy medium. (/ml culture/5ml ponassy).
2. Incubate it for 2 hrs. at 37°C. (To make new culture.)
4. Mix them in the ratio of 1 : 10^6 cells/ml

No of tube	1	2	3	4	control
W6 F8 :	$\times 10^{-2}$ ↓ 2×10^{6}	10^{-2} ↓ 2×10^{4}	$\times 10^{-2}$ ↓ 2×10^{2}	$\times 10^{-1}$ ↓ 2×10^{1}	~~+~~ ⇐ 0
	0.1ml ↓ 2×10^{5} +	0.1ml ↓ 2×10^{3} +	0.1ml ↓ 2×10^{1} +	0.1ml ↓ 2×10^{0} +	~~+~~ 0
W3086 Ca. 10^8	1ml	1ml	1ml	1ml	1ml.

5. Add 5m into the mixture. at the final concentration of ~~1000~~ 1,000 unit /ml. to kill F8 donor. ~~after~~ 2 hrs after mixing, and incubate it overnight. 4:00 pm. 5/V/59.

2g/5ml H_2O $\times 10^{-1}$: 0.2ml/1ml pen. final : 0.8 mg /ml.

6. Survival count of W6 F8. & W3086.

W6 F8: $\times10^{-2}\times10^{-2}\times10^{-1}\times10^{-1}$ (0.1ml/plate) — 230, 189

W3086: $\times10^{-2}\times10^{-2}\times10^{-1}\times10^{-1}$ 0.1ml. — 206, 185

Cell Numbers used : Original cell #. : 2 × 10⁸ cells/ml : 2 × 10⁸ cells/ml.

number of F8 donor

W6 F8	3086
# of F'	# of F'
2×10^5	10^8
2×10^3	10^8
2×10^1	10^8
2×10^0	10^8
~~0~~	10^8
W6 F8	—



Summary of experimental conditions.

young culture : ~~→~~ 2 hrs. culture in pon. at 37°C.

Inoculum size :

Minimal cell number for infection of F' (within 2 hrs.)

$2\times10^3 \sim 2\times10^1$

$2000 \sim 20$

Ca. 100^2.

F4 → F8.

×2979 on MGal

④ type.



on MXyl. ×2979.

F4 → F8.

×2979 on MGal

⑧ type

on MXyl ×2979.

Segregation pattern of F_8 and F_4 from double F. ($F_{4,8}$). judged by Replica plating method

cf. ♯42

Compair these two segregant. above 4 type ; bellow 8 type.
Xyl plate is ok : F4 type is more futile on MXyl, than F8 type.
But MGal plate is not suitable for distination between F8 and F4.

Use #2 and #31. to detect recombinats between F'.