

Reprinted from SCIENCE, April 12, 1957, Vol. 125, No. 3250, pages 691-692.

## Reductive Cleavage of Disulfide Bridges in Ribonuclease

Michael Sela, Frederick H. White, Jr., and Christian B. Anfinsen  
*Laboratory of Cellular Physiology and Metabolism, National Heart Institute,  
National Institutes of Health, Bethesda, Maryland*

## Reductive Cleavage of Disulfide Bridges in Ribonuclease

Total cleavage of the four disulfide bridges in ribonuclease (RNase) may be achieved with complete loss of enzyme activity, as assayed (1) with ribonucleic acid as substrate. Bovine pancreatic ribonuclease (Armour) (12 mg/ml) was reacted with thioglycolic acid [400 moles per mole of enzyme, assuming 13,683 as the molecular weight of ribonuclease, (2) and a water content of 9.38 percent in the air-dried preparation] in 8*M* urea at pH 8.5 (adjusted with aqueous trimethylamine) for 4.5 hours at room temperature. The reduced ribonuclease was precipitated with acetone and 1*M* hydrochloric acid (39/1) at -5°C, and the sample was washed three times with this solvent and twice with cold ether. The material so obtained was completely reduced, as indicated by the presence of eight sulfhydryl groups per mole of enzyme on spectrophotometric titration with *p*-chloromercuribenzoate (3).

An alternative method that was used for following the extent of reduction involved selective carboxymethylation of the sulfhydryl groups of reduced ribonuclease (14 mg/ml) by reaction, under nitrogen, with iodoacetic acid (700 moles per mole of protein) for 2 hours at room temperature in an autotitrator. Trimethylamine (5 percent aqueous solution) was added continuously to maintain the pH at 8.5. The reaction was stopped, and the carboxymethylated protein was precipitated and washed as described for the reduced material. (The carboxymethylated protein was used as a test for the completeness of this washing procedure by its treatment with thioglycollate as in the reduction procedure, with subsequent washing as described. Titration with *p*-chloromercuribenzoate of the protein so treated revealed no sulfhydryl groups, thus establishing the complete removal of thioglycollate from the reduced enzyme by

the washing procedure.) The product was hydrolyzed for 16 to 18 hours in 6*N* hydrochloric acid at 100°C. S-Carboxymethylcysteine is known to be stable under these conditions of hydrolysis (4). The hydrolyzate was dinitrophenylated (5) and chromatographed in two dimensions. A mixture of toluene, pyridine, chloroethanol, and 0.8*N* ammonium hydroxide (10/3/6/6) was used in the first dimension (5, 6), and, in the second, tertiary amyl alcohol and 0.2*M* phthalate buffer at pH 5 (2/1) (7). The component which moved identically with an authentic sample of dinitrophenyl-S-carboxymethylcysteine was eluted, and its concentration was determined spectrophotometrically (5). The chromatographic analysis revealed 8 moles of dinitrophenyl-S-carboxymethylcysteine per mole of reduced, carboxymethylated ribonuclease, but no trace of bis-dinitrophenylcystine. This method of analysis thus confirmed the completeness of cleavage of disulfide bridges as determined by titration with *p*-chloromercuribenzoate.

Carrying out the reductions in 8*M* urea for shorter periods of time, with a smaller excess of thioglycollate, or at slightly lower pH values, yielded partially reduced materials with varying degrees of activity (see Fig. 1).

In the absence of urea, but with other conditions identical with those described for the reduction of ribonuclease, complete cleavage of disulfide bridges in the enzyme was not achieved. This finding parallels that of Lindley (8), who demonstrated that only one of the three disulfide bridges in insulin was reductively cleaved by thioglycolic acid in the absence of urea. In the present experiments, production of sulfhydryl groups was linear up to 30 minutes, reaching a value of 3.7. After 2 hours, 4.6 sulfhydryl groups were formed. It appears that two of the disulfide bridges in ribonuclease may be opened with relative ease, but that the remaining two are cleaved appreciably only in the presence of urea. In contrast to the results experienced with complete reduction, enzyme activity was diminished but not entirely lost during partial reduction. A correlation between the number of sulfhydryl groups per mole and enzyme activity is given in Fig. 1, which summarizes all reduction experiments performed. Carboxymethylation of the partially reduced material caused no further reduction in enzyme activity, nor did the conditions of car-

boxymethylation affect the activity of native ribonuclease.

When completely reduced and fully inactive enzyme (1 mg/ml) was subjected to oxidation by air bubbling at room temperature for 68 hours in 0.01*M* phosphate buffer at pH 7 to 8, ribonuclease activity reappeared to the extent of from 12 to 19 percent of the specific activity of native enzyme. The number of sulfhydryl groups per mole decreased during reoxidation. When partially reduced ribonuclease containing about four sulfhydryl groups per mole was reoxidized, there was no significant increase in activity or decrease in sulfhydryl groups. The results of all reoxidation experiments are summarized in Fig. 1. Native ribonuclease that was treated under these reoxidation conditions showed no change in activity, nor was activity regenerated in completely reduced, carboxymethylated ribonuclease under these conditions.

The disappearance of disulfide bridges from native ribonuclease during reduction, and their reappearance on oxidation of reduced enzyme, was followed qualitatively by subtilisin digestion (9) of the carboxymethylated protein and




Fig. 1. Activity of ribonuclease at various stages of reduction (expressed as percentage of the specific activity of native ribonuclease) as a function of the number of moles of sulfhydryl per mole of enzyme. ▲, Reduction in absence of urea; ●, reduction in 8*M* urea; □, reoxidation of fully reduced, inactive ribonuclease; ○, reoxidation of samples containing more than six sulfhydryl groups per average molecule; ▽, reoxidation of samples containing about four sulfhydryl groups per average molecule.

All technical papers and comments on them are published in this section. Manuscripts should be typed double-spaced and be submitted in duplicate. In length, they should be limited to the equivalent of 1200 words; this includes the space occupied by illustrative or tabular material, references and notes, and the author(s)' name(s) and affiliation(s). Illustrative material should be limited to one table or one figure. All explanatory notes, including acknowledgments and authorization for publication, and literature references are to be numbered consecutively, keyed into the text proper, and placed at the end of the article under the heading "References and Notes." For fuller details see "Suggestions to Contributors" in *Science* 125, 16 (4 Jan. 1957).

paper electrophoresis (10) of the digest. Native ribonuclease treated in this way (11) yields several bands which contain disulfide bonds as shown by the cyanide-nitroprusside test (12). The stepwise disappearance of these bands on reduction and their reappearance on oxidation of reduced ribonuclease have been observed. Identification of the disulfide-containing bands resulting from the digestion of reoxidized preparations with those produced from native ribonuclease, as well as elucidation of the order of cleavage of the disulfide bonds during the reduction of the native molecule, must await further investigation.

The S-shaped distribution of the solid points in Fig. 1 suggests the formation of enzymatically active products at intermediate stages of reduction. Further, this figure indicates that, with as many as four sulfhydryl groups per average molecule, activity remains high. With the appearance of more than this number of sulfhydryl groups, activity drops

rapidly. Positive confirmation of the existence of enzymatically active reduction intermediates will depend on their separation and characterization.

Various covalent (9, 13), secondary (14, 15) and tertiary (15) bonds can be broken in ribonuclease without loss of enzymatic activity, thus suggesting that the "active center" of the enzyme might comprise only a relatively small part of the molecule (14, 16). It appears from the present work that not all of the disulfide bridges in ribonuclease are essential for enzyme activity. A further study of the intermediate stages of ribonuclease reduction may reveal specific disulfide bridges associated with the "active center" of this enzyme.

MICHAEL SELA\*

FREDERICK H. WHITE, JR.

CHRISTIAN B. ANFINSEN

*Laboratory of Cellular Physiology  
and Metabolism, National Heart  
Institute, National Institutes of  
Health, Bethesda, Maryland*

#### References and Notes

1. C. B. Anfinsen *et al.*, *J. Biol. Chem.* 207, 201 (1954).
  2. C. H. W. Hirs, W. H. Stein, S. Moore, *ibid.* 219, 623 (1956).
  3. P. D. Boyer, *J. Am. Chem. Soc.* 76, 4331 (1954).
  4. A. Berger, J. Noguchi, E. Katchalski, *ibid.* 78, 4483 (1956).
  5. A. L. Levy, *Nature* 174, 126 (1954).
  6. G. Biserte and R. Osteux, *Bull. soc. chim. biol.* 33, 50 (1951).
  7. S. Blackburn and A. G. Lowther, *Biochem. J. (London)* 48, 126 (1951).
  8. H. Lindley, *J. Am. Chem. Soc.* 77, 4927 (1955).
  9. F. M. Richards, *Compt. rend. trav. lab. Carlsberg Sér. chim.* 29, 329 (1955).
  10. H. Michl, *Monatsh. Chem.* 82, 489 (1951).
  11. C. B. Anfinsen and A. Ryle, in preparation.
  12. G. Toennies and J. J. Kolb, *Anal. Chem.* 23, 823 (1951).
  13. G. Kalnitsky and W. J. Rogers, *Biochim. et Biophys. Acta* 20, 378 (1956).
  14. C. B. Anfinsen, *Compt. rend. trav. lab. Carlsberg Sér. chim.* 30, 13 (1956).
  15. M. Sela and C. B. Anfinsen, *Biochim. et Biophys. Acta*, in press.
  16. C. B. Anfinsen and R. R. Redfield, *Advances in Protein Chem.* 11, 1 (1956).
- \* On leave of absence from the Weizmann Institute of Science, Rehovoth, Israel.

1 February 1957