

THE HEALTH CONSEQUENCES OF INVOLUNTARY SMOKING

a report of the Surgeon General

1986



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
Centers for Disease Control
Center for Health Promotion and Education
Office on Smoking and Health
Rockville, Maryland 20857



THE SECRETARY OF HEALTH AND HUMAN SERVICES
WASHINGTON, D.C. 20201

DEC 15 1986

The Honorable George Bush
President of the Senate
Washington, D.C. 20510

Dear Mr. President:

It is my pleasure to transmit to the Congress the 1986 Surgeon General's Report on the health consequences of smoking, as mandated by Section 8(a) of the Public Health Cigarette Smoking Act of 1969. The current volume, entitled The Health Consequences of Involuntary Smoking, examines the scientific evidence on the health effects resulting from nonsmoker exposure to environmental tobacco smoke.

The issue of whether or not tobacco smoke is carcinogenic for humans was conclusively resolved more than 20 years ago when the first report on smoking and health was issued in 1964. Based on the current report, the judgment can now be made that exposure to environmental tobacco smoke can cause disease, including lung cancer, in nonsmokers. It is also clear that simple separation of smokers and nonsmokers within the same airspace may reduce but cannot eliminate nonsmoker exposure to environmental tobacco smoke.

The report also reviews an extensive body of evidence which establishes an increased risk of respiratory illness and reduced lung function in infants and very young children of parents who smoke. This effect is more pronounced if both parents smoke than if only one parent smokes. As a physician, I believe that parents should refrain from smoking around small children both as a means of protecting their children's health and to set a good example for the child.

Today, only 30 percent of the adult population in the United States are smokers--the lowest level of smoking in the country since World War II, reflecting that the great majority of the population has never smoked or has successfully quit.

Accompanying this decline in overall prevalence of cigarette smoking has been an increased concern for protecting the health and well being of nonsmokers, as evidenced by the number of laws and regulations restricting smoking in public places. Today, 40 States and the District of Columbia have enacted some form of legislation to restrict smoking in public. Increasingly, these laws pertain to protecting nonsmokers in many different settings, including the workplace.

Based on the evidence presented in this report, the choice to smoke should not interfere with the nonsmoker's choice for an environment free of tobacco smoke.

Sincerely,

Otis R. Bowen, M.D.
Secretary

Enclosure



THE SECRETARY OF HEALTH AND HUMAN SERVICES
WASHINGTON, D.C. 20201

DEC 5

The Honorable Thomas P. O'Neill, Jr.
Speaker of the House
of Representatives
Washington, D.C. 20515

Dear Mr. Speaker:

It is my pleasure to transmit to the Congress the 1986 Surgeon General's Report on the health consequences of smoking, as mandated by Section 8(a) of the Public Health Cigarette Smoking Act of 1969. The current volume, entitled The Health Consequences of Involuntary Smoking, examines the scientific evidence on the health effects resulting from nonsmoker exposure to environmental tobacco smoke.

The issue of whether or not tobacco smoke is carcinogenic for humans was conclusively resolved more than 20 years ago when the first report on smoking and health was issued in 1964. Based on the current report, the judgment can now be made that exposure to environmental tobacco smoke can cause disease, including lung cancer, in nonsmokers. It is also clear that simple separation of smokers and nonsmokers within the same airspace may reduce but cannot eliminate nonsmoker exposure to environmental tobacco smoke.

The report also reviews an extensive body of evidence which establishes an increased risk of respiratory illness and reduced lung function in infants and very young children of parents who smoke. This effect is more pronounced if both parents smoke than if only one parent smokes. As a physician, I believe that parents should refrain from smoking around small children both as a means of protecting their children's health and to set a good example for the child.

Today, only 30 percent of the adult population in the United States are smokers—the lowest level of smoking in the country since World War II, reflecting that the great majority of the population has never smoked or has successfully quit.

Accompanying this decline in overall prevalence of cigarette smoking has been an increased concern for protecting the health and well being of nonsmokers, as evidenced by the number of laws and regulations restricting smoking in public places. Today, 40 States and the District of Columbia have enacted some form of legislation to restrict smoking in public. Increasingly, these laws pertain to protecting nonsmokers in many different settings, including the workplace.

Based on the evidence presented in this report, the choice to smoke should not interfere with the nonsmoker's choice for an environment free of tobacco smoke.

Sincerely,

Otis R. Bowen, M.D.
Secretary

Enclosure

FOREWORD

The data reviewed in 17 previous U.S. Public Health Service reports on the health consequences of smoking have conclusively established cigarette smoking as the largest single preventable cause of premature death and disability in the United States.

The question whether tobacco smoke is harmful to smokers was answered more than 20 years ago. As a result, many scientists began to question whether the low levels of exposure to environmental tobacco smoke (ETS) received by nonsmokers could also be harmful.

The current Report, *The Health Consequences of Involuntary Smoking*, examines the evidence that even the lower exposure to smoke received by the nonsmoker carries with it a health risk. Use of the term "involuntary smoking" denotes that for many nonsmokers, exposure to ETS is the result of an unavoidable consequence of being in proximity to smokers. It is the first Report in the health consequences of smoking series to establish a health risk due to tobacco smoke exposure for individuals other than the smoker, and represents the work of more than 60 distinguished physicians and scientists, both in this country and abroad.

After careful examination of the available evidence, the following overall conclusions can be reached:

1. Involuntary smoking is a cause of disease, including lung cancer, in healthy nonsmokers.
2. The children of parents who smoke, compared with the children of nonsmoking parents, have an increased frequency of respiratory infections, increased respiratory symptoms, and slightly smaller rates of increase in lung function as the lung matures.
3. Simple separation of smokers and nonsmokers within the same air space may reduce, but does not eliminate, exposure of nonsmokers to environmental tobacco smoke.

Exposure to environmental tobacco smoke occurs at home, at the worksite, in public, and in other places where smoking is permitted.

The quality of the indoor environment must be a concern of all who control and occupy that environment. Protection of individuals from exposure to environmental tobacco smoke is therefore a responsibility shared by all:

- As parents and adults we must protect the health of our children by not exposing them to environmental tobacco smoke.
- As employers and employees we must ensure that the act of smoking does not expose the nonsmoker to tobacco smoke.
- For smokers, it is their responsibility to assure that their behavior does not jeopardize the health of others.
- For nonsmokers, it is their responsibility to provide a supportive environment for smokers who are attempting to stop.

Actions taken by individuals, employers, and employee organizations reflect the growing concern for protecting nonsmokers. The number of laws and regulations enacted at the national, State, and local level governing smoking in public has increased substantially over the past 10 years, and surveys conducted by numerous organizations show strong public support for these actions among both smokers and nonsmokers.

As a Nation, we have made substantial progress in addressing the enormous toll inflicted by active smoking. Efforts to improve and protect individual health must be not only continued but strengthened. On the basis of the evidence presented in this Report, it is clear that actions to protect nonsmokers from ETS exposure not only are warranted but are essential to protect public health.

Robert E. Windom, M.D.
Assistant Secretary for Health

PREFACE

This, the 1986 Report of the Surgeon General, is the U.S. Public Health Service's 18th in the health consequences of smoking series and the 5th issued during my tenure as Surgeon General.

Previous Reports have documented the tremendous health burden to society from smoking, particularly cigarette smoking. The evidence establishing cigarette smoking as the single largest preventable cause of premature death and disability in the United States is overwhelming—totaling more than 50,000 studies from dozens of cultures. Smoking is now known to be causally related to a variety of cancers in addition to lung cancer; it is a cause of cardiovascular disease, particularly coronary heart disease, and is the major cause of chronic obstructive lung disease. It is estimated that smoking is responsible for well over 300,000 deaths annually in the United States, representing approximately 15 percent of all mortality.

Thirty years ago, however, the scientific evidence linking smoking with early death and disability was more limited. By 1964, the year the Advisory Committee to the Surgeon General issued the first report on smoking and health, a substantial body of evidence had accumulated upon which a judgment could be made that smoking was a cause of disease in active smokers. Subsequent reports over the last 20 years have expanded our understanding and knowledge about smoking behavior, the toxicity and carcinogenicity of tobacco smoke, and the specific disease risks resulting from exposure to this agent.

This Report is the first issued since 1964 that identifies a chronic disease risk resulting from exposure to tobacco smoke for individuals other than smokers. It is now clear that disease risk due to the inhalation of tobacco smoke is not limited to the individual who is smoking, but can extend to those who inhale tobacco smoke emitted into the air. This Report represents a detailed review of the health effects resulting from nonsmoker exposure to environmental tobacco smoke (ETS). ETS is the combination of smoke emitted from a burning tobacco product between puffs (sidestream smoke) and the smoke exhaled by the smoker. The 1986 Report, *The Health Consequences of Involuntary Smoking*, is a critical review of all the available scientific evidence pertaining to the health effects of ETS exposure on nonsmokers. The term "involuntary smoking" is used to

note that such exposures often occur as an unavoidable consequence of being in close proximity to smokers.

Lung Cancer and Environmental Tobacco Smoke

The appropriate framework for an examination of the lung cancer risk from involuntary smoking is that of a low-dose exposure to a known human carcinogen. Over 30 years of research have conclusively established cigarette smoke as a carcinogen. This Report presents evidence that the chemical composition of sidestream smoke is qualitatively similar to the mainstream smoke inhaled by the active smoker, and that both mainstream and sidestream smoke act as carcinogens in bioassay systems. Data related to environmental levels of tobacco smoke constituents and from measures of nicotine absorption in nonsmokers suggest that nonsmokers are exposed to levels of environmental tobacco smoke that would be expected to generate a lung cancer risk; epidemiological studies of populations exposed to ETS have documented an increased risk for lung cancer in those nonsmokers with increased exposure.

It is rare to have such detailed exposure data or human epidemiologic studies on disease occurrence when attempting to evaluate the risk of low-dose exposure to an agent with established toxicity at higher levels of exposure. The relative abundance of data reviewed in this Report, their cohesiveness, and their biologic plausibility allow a judgment that involuntary smoking can cause lung cancer in nonsmokers. Although the number of lung cancers due to involuntary smoking is smaller than that due to active smoking, it still represents a number sufficiently large to generate substantial public health concern.

It is certain that a substantial proportion of the lung cancers that occur in nonsmokers are due to ETS exposure; however, more complete data on the dose and variability of smoke exposure in the nonsmoking U.S. population will be needed before a quantitative estimate of the number of such cancers can be made.

Children and Infants

This Report also documents a relationship between parental smoking and the respiratory health of infants and children (under 2 years of age). Infants of parents who smoke have an increased risk of hospitalization for bronchitis and pneumonia when compared with infants of nonsmoking parents. There is a relationship between parental smoking and an increased frequency of respiratory symptoms in children. A slower rate of growth in lung function has been observed in children of smoking parents. In many studies, if both

parents smoke, a stronger relationship exists than if only one parent smokes.

What future respiratory burden these findings may represent for these children later in life is not known. As a former pediatric surgeon, I strongly urge parents to refrain from smoking in the presence of children as a means of protecting not only their children's current health status but also their own.

Diseases Other Than Lung Cancer

Several studies have provided data on the relationship between ETS and cancers other than lung cancer and on ETS exposure and cardiovascular disease. However, further research in these areas will be required to determine whether an association exists between ETS exposure and an increased risk of developing these diseases.

Policies Restricting Smoking in Public Places

The growth in our understanding of the disease risk associated with involuntary smoking has been accompanied by a change in the social acceptability of smoking and by a growing body of legislation, regulation, and voluntary action that addresses where smoking may occur in public. Forty States and the District of Columbia now have some form of legislation controlling or restricting smoking in various public settings. Some States limit smoking to only a few designated areas; however, States are increasingly developing and implementing comprehensive legislation that restricts smoking in many public settings, including the workplace. Nine States have restrictions that cover smoking not only by public employees but also by employees in the private sector.

No systematic evaluation of the effects these measures may have on smoking behavior has been conducted, but there is little doubt that strong public sentiment exists for implementing such restrictions. A number of national surveys conducted by voluntary health organizations, government agencies, and even the tobacco industry have documented that an overwhelming majority of both smokers and nonsmokers support restricting smoking in public.

Public Health Policy and Involuntary Smoking

The 1986 Surgeon General's Report on the Health Consequences of Involuntary Smoking clearly documents that nonsmokers are placed at increased risk for developing disease as the result of exposure to environmental tobacco smoke.

Critics often express that more research is required, that certain studies are flawed, or that we should delay action until more conclusive proof is produced. As both a physician and a public health

official, it is my judgment that the time for delay is past; measures to protect the public health are required now. The scientific case against involuntary smoking as a health risk is more than sufficient to justify appropriate remedial action, and the goal of any remedial action must be to protect the nonsmoker from environmental tobacco smoke.

The data contained in this Report on the rapid diffusion of tobacco smoke throughout an enclosed environment suggest that separation of smokers and nonsmokers in the same room or in different rooms that share the same ventilation system may reduce ETS exposure but will not eliminate exposure. The responsibility to protect the safety of the indoor environment is shared by all who occupy or control that environment.

Changes in smoking policies regarding the workplace and other environments necessitated by the data presented in this Report should not be designed to punish the smoker. Successful implementation of protection for the nonsmoker requires the support and cooperation of smokers, nonsmokers, management, and employees and should be developed through a cooperative effort of all groups affected. In addition, changes are often more effective when support and assistance is provided for the smoker who wants to quit.

Cigarette smoking is an addictive behavior, and the individual smoker must decide whether or not to continue that behavior; however, it is evident from the data presented in this volume that the choice to smoke cannot interfere with the nonsmokers' right to breathe air free of tobacco smoke. The right of smokers to smoke ends where their behavior affects the health and well-being of others; furthermore, it is the smokers' responsibility to ensure that they do not expose nonsmokers to the potential harmful effects of tobacco smoke.

C. Everett Koop, M.D.
Surgeon General

ACKNOWLEDGMENTS

This Report was prepared by the Department of Health and Human Services under the general editorship of the Office on Smoking and Health, Donald R. Shopland, Acting Director. Managing Editor was William R. Lynn, Acting Technical Information Officer, Office on Smoking and Health.

Senior scientific editor was David M. Burns, M.D., Associate Professor of Medicine, Division of Pulmonary and Critical Care Medicine, University of California Medical Center, San Diego, San Diego, California. Consulting scientific editors were Ellen R. Gritz, Ph.D., Director, Division of Cancer Control, Jonsson Comprehensive Cancer Center, University of California, Los Angeles, Los Angeles, California; John H. Holbrook, M.D., Associate Professor of Internal Medicine, Department of Internal Medicine, University Hospital, Salt Lake City, Utah; and Jonathan M. Samet, M.D., Professor of Medicine, Department of Medicine, The University of New Mexico School of Medicine, Albuquerque, New Mexico.

The following individuals prepared draft chapters or portions of the Report.

Neal Benowitz, M.D., San Francisco General Medical Center, San Francisco, California

A. Sonia Buist, M.D., Professor of Medicine, Department of Physiology, Oregon Health Sciences University, Portland, Oregon

Charles Hiller, M.D., Pulmonary Division, University Hospital, Little Rock, Arkansas

Dietrich Hoffmann, Ph.D., Associate Director, Naylor Dana Institute for Disease Prevention, American Health Foundation, Valhalla, New York

Ilse Hoffmann, Research Coordinator, Naylor Dana Institute for Disease Prevention, American Health Foundation, Valhalla, New York

John R. Hoidal, M.D., Director of Pulmonary Medicine, University of Tennessee Center for Health Sciences, Memphis, Tennessee

John McCarthy, M.P.H., Harvard School of Public Health, Boston, Massachusetts

Nancy A. Rigotti, M.D., Institute for the Study of Smoking Behavior and Policy, John F. Kennedy School of Government, Harvard University, Cambridge, Massachusetts

Jonathan M. Samet, M.D., Professor of Medicine, Department of Medicine, The University of New Mexico School of Medicine, Albuquerque, New Mexico

John Spengler, Ph.D., Harvard School of Public Health, Boston, Massachusetts

Annetta Weber, Ph.D., Federal Institute of Technology, Zurich, Switzerland

Scott T. Weiss, M.D., M.S., Associate Professor of Medicine, Channing Laboratories, Harvard Medical School, Boston, Massachusetts

Anna H. Wu, Ph.D., Department of Preventive Medicine, School of Medicine, University of Southern California, Los Angeles, California

The editors acknowledge with gratitude the following distinguished scientists, physicians, and others who lent their support in the development of this Report by coordinating manuscript preparation, contributing critical reviews of the manuscript, or assisting in other ways.

Elvin E. Adams, M.D., M.P.H., Director, Health and Temperance Department, General Conference of Seventh-Day Adventists, Washington, D.C.

Stephen M. Ayres, M.D., Dean, School of Medicine, Medical College of Virginia, Richmond, Virginia

David V. Bates, M.D., Professor of Medicine and Physiology, Department of Medicine, Acute Care Hospital, University of British Columbia, Vancouver, British Columbia

William J. Blot, Ph.D., Chief, Biostatistics Branch, Epidemiology and Biostatistics Program, Division of Etiology, National Cancer Institute, National Institutes of Health, Bethesda, Maryland

Benjamin Burrows, M.D., Professor of Internal Medicine, and Director, Division of Respiratory Sciences, The University of Arizona College of Medicine, Tucson, Arizona

D. M. DeMarini, Ph.D., Genetic Toxicology Division, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina

Vincent T. DeVita, Jr., M.D., Director, National Cancer Institute, National Institutes of Health, Bethesda, Maryland

Louis Diamond, Ph.D., College of Pharmacy, University of Kentucky, Lexington, Kentucky

Richard Doll, Cancer Epidemiology and Clinical Trials Unit, Imperial Cancer Research Fund, The Radcliffe Infirmary, University of Oxford, Oxford, England, United Kingdom

Manning Feinleib, M.D., Dr.P.H., Director, National Center for Health Statistics, Office of the Assistant Secretary for Health, Hyattsville, Maryland

Edwin B. Fisher, Jr., Ph.D., Associate Professor, Department of Psychology, Washington University, St. Louis, Missouri

William H. Foege, M.D., Executive Director, Task Force for Child Survival, Carter Presidential Center, Atlanta, Georgia

Joseph F. Fraumeni, Jr., M.D., Associate Director for Epidemiology and Biostatistics, Division of Cancer Etiology, National Cancer Institute, National Institutes of Health, Bethesda, Maryland

Lawrence Garfinkel, M.A., Vice President for Epidemiology and Statistics, and Director of Cancer Prevention, American Cancer Society, New York, New York

R.A. Griesemer, D.V.M., Ph.D., Director, Biology Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee

Michael R. Guerin, Ph.D., Organic Chemistry Section, Analytical Chemistry, Oak Ridge National Laboratory, Oak Ridge, Tennessee

Jeffery E. Harris, M.D., Ph.D., Associate Professor, Department of Economics, Massachusetts Institute of Technology, Cambridge, Massachusetts

Millicent Higgins, M.D., Associate Director, Epidemiology and Biometry Program, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, Maryland

Takeshi Hirayama, M.D., Director, Institute of Preventive Oncology, Shinjuku-ku, Tokyo, Japan

Dwight Janerich, D.D.S., M.P.H., Department of Epidemiology and Public Health, Yale University School of Medicine, New Haven, Connecticut

Martin Jarvis, M.P.H., Senior Clinical Psychologist, Addiction Research Unit, Institute of Psychiatry, London, England, United Kingdom

Brian P. Leaderer, Ph.D., M.P.H., Associate Fellow, John B. Pierce Foundation Laboratory, Associate Professor, Department of Epidemiology and Public Health, Yale University School of Medicine, New Haven, Connecticut

Charles L. LeMaistre, M.D., President, University of Texas Systems Cancer Center, Houston, Texas

Claude Lenfant, M.D., Director, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, Maryland

Donald Ian Macdonald, M.D., Administrator, Alcohol, Drug Abuse, and Mental Health Administration, Rockville, Maryland

James S. Marks, M.D., M.P.H., Assistant Director for Science, Center for Health Promotion and Education, Centers for Disease Control, Atlanta, Georgia

James O. Mason, M.D., Dr.P.H., Director, Centers for Disease Control, Atlanta, Georgia

J. Michael McGinnis, M.D., Deputy Assistant Secretary for Health (Disease Prevention and Health Promotion), Office of the Assistant Secretary for Health, Washington, D.C.

A. J. McMichael, M.D., M.B.B.S., Ph.D., Chairman and Senior Principal Research Scientist, CSIRO Division of Human Nutrition, Adelaide, South Australia

D. J. Moschandreas, Ph.D., Research Director, ITT Research Institute, Chicago, Illinois

David Muir, M.D., Director, Occupational Health Program, Health Sciences Center, McMaster University, Hamilton, Ontario, Canada

C. Tracy Orleans, Ph.D., Research Associate, Health Services Research Center, University of North Carolina, Chapel Hill, North Carolina

Richard Peto, M.A., M.Sc., I.C.R.F., Regius Assessor of Medicine, The Radcliffe Infirmary, University of Oxford, Oxford, England, United Kingdom

Otto Raabe, M.D., Laboratory for Energy Related Health Research, University of California, Davis, Davis, California

James L. Repace, Chief of Technical Services, Indoor Air Quality Program, U.S. Environmental Protection Agency, Washington, D.C.

M.A.H. Russell, F.R.C.P., Addiction Research Unit, Institute of Psychiatry, University of London, London, England, United Kingdom

Roy J. Shephard, M.D., Ph.D., Director, School of Physical and Health Education, University of Toronto, Toronto, Canada

Frank E. Speizer, M.D., Channing Laboratories, Harvard Medical School, Boston, Massachusetts

Jesse L. Steinfeld, M.D., President, Medical College of Georgia, Augusta, Georgia

David N. Sundwall, M.D., Administrator, Health Resources and Services Administration, Rockville, Maryland

Gregory W. Traynor, Staff Scientist, Lawrence Berkeley Laboratory, Berkeley, California

Dimitrios Trichopoulos, Director, Department of Hygiene and Epidemiology, School of Medicine, University of Athens, Athens, Greece

Kenneth E. Warner, Ph.D., Professor, and Chairman, Department of Public Health Policy and Administration, School of Public Health, The University of Michigan, Ann Arbor, Michigan

Ernst L. Wynder, M.D., President, American Health Foundation, New York, New York

James B. Wyngaarden, M.D., Director, National Institutes of Health, Bethesda, Maryland

Frank E. Young, M.D., Commissioner, Food and Drug Administration, Rockville, Maryland

The editors also acknowledge the contributions of the following staff members and others who assisted in the preparation of this Report.

Erica W. Adams, Chief Copy Editor and Assistant Production Manager, Health and Natural Resources Department, Sterling Software, Inc., Rockville, Maryland

Richard H. Amacher, Director, Health and Natural Resources Department, Sterling Software, Inc., Rockville, Maryland

Margaret L. Anglin, Secretary, Office on Smoking and Health, Rockville, Maryland

John L. Bagrosky, Associate Director for Program Operations, Office on Smoking and Health, Rockville, Maryland

Charles A. Brown, Programmer, Automation and Technical Services Department, Sterling Software, Inc., Rockville, Maryland

Clarice D. Brown, Statistician, Office on Smoking and Health, Rockville, Maryland

Richard C. Brubaker, Information Specialist, Health and Natural Resources Department, Sterling Software, Inc., Rockville, Maryland

Catherine E. Burckhardt, Secretary, Office on Smoking and Health, Rockville, Maryland

Joanna B. Crichton, Copy Editor, Health and Natural Resources Department, Sterling Software, Inc., Rockville, Maryland

Stephanie D. DeVoe, Programmer, Automation and Technical Services Department, Sterling Software, Inc., Rockville, Maryland

Danny A. Goodman, Information Specialist, Health and Natural Resources Department, Sterling Software, Inc., Rockville, Maryland

Patricia E. Healy, Technical Information Specialist, Office on Smoking and Health, Rockville, Maryland

Terri L. Henry, Clerk-Typist, Office on Smoking and Health, Rockville, Maryland

Timothy K. Hensley, Technical Publications Writer, Office on Smoking and Health, Rockville, Maryland

Shirley K. Hickman, Data Entry Operator, Health and Natural Resources Department, Sterling Software, Inc., Rockville, Maryland

Robert S. Hutchings, Associate Director for Information and Program Development, Office on Smoking and Health, Rockville, Maryland

Maureen Illar, Editorial Assistant, Office on Smoking and Health, Rockville, Maryland

Julie Kurz, Graphic Artist, Information Center Management Department, Sterling Software, Inc., Rockville, Maryland

Ruth C. Palmer, Secretary, Office on Smoking and Health, Rockville, Maryland

Jerome A. Paulson, M.D., Medical Officer, Office on Smoking and Health, Rockville, Maryland

Russell D. Peek, Library Acquisitions Specialist, Health and Natural Resources Department, Sterling Software, Inc., Rockville, Maryland

Margaret E. Pickerel, Public Information and Publications Specialist, Office on Smoking and Health, Rockville, Maryland

Raymond K. Poole, Production Coordinator, Health and Natural Resources Department, Sterling Software, Inc., Rockville, Maryland

Linda R. Spiegelman, Administrative Officer, Office on Smoking and Health, Rockville, Maryland

Evelyn L. Swarr, Administrative Secretary, Automation and Technical Services Department, Sterling Software, Inc., Rockville, Maryland

Debra C. Tate, Publications Systems Specialist, Publishing Systems Division, Sterling Software, Inc., Riverdale, Maryland

Jerry W. Vaughn, Programmer, University of California, San Diego, San Diego, California

Mary I. Walz, Computer Systems Analyst, Office on Smoking and Health, Rockville, Maryland

Louise G. Wiseman, Technical Information Specialist, Office on Smoking and Health, Rockville, Maryland

Pamela Zuniga, Secretary, University of California, San Diego, San Diego, California

TABLE OF CONTENTS

Foreword	vii
Preface.....	ix
Acknowledgments.....	xiii
1. Introduction, Overview, and Summary and Conclusions	1
2. Health Effects of Environmental Tobacco Smoke Ex- posure.....	17
3. Environmental Tobacco Smoke Chemistry and Expo- sures of Nonsmokers	121
4. Deposition and Absorption of Tobacco Smoke Con- stituents	177
5. Toxicity, Acute Irritant Effects, and Carcinogenicity of Environmental Tobacco Smoke	225
6. Policies Restricting Smoking in Public Places and the Workplace.....	261
Index	335
